

AEM012-P02

Room: Convention Hall

Time: May 27 17:15-18:45

## Atmospheric electric field and ELF observation for global-circuit model

Ryoe Sato<sup>1\*</sup>, Masashi Kamogawa<sup>1</sup>, Yukihiro Takahashi<sup>2</sup>, Hironobu Fujiwara<sup>3</sup>, Isao Yamamoto<sup>4</sup>, Mitsuteru Sato<sup>5</sup>, Yoshihiro Kakinami<sup>6</sup>, Yoshihisa Kimura<sup>1</sup>

<sup>1</sup>Dpt. of Phys., Tokyo Gakugei Univ., <sup>2</sup>Dpt. Cosmosci., Hokkaido Univ., <sup>3</sup>Joshiseigakuin High School,

<sup>4</sup>Dpt. of Info. Comp. Eng., Okayama Univ., <sup>5</sup>Dpt. Cosmosci., Hokkaido Univ.,

<sup>6</sup>Institute of Space Sci., National Centra

In early 20th century, global circuit (GC) model was proposed. After the observation of Carnegie research ship, the concept of the global electrical circuit was established. Although many applied models have been proposed, the initial GC model has been practically used to understand global atmospheric electric field (AEF) variations. However, it is very difficult to extract the global component of AEF, because AEF is highly affected by ion-aerosol variation in fair weather especially in the continent. Therefore, most of extracting the global component is operated by taking the average of AEF time-series in the fair weather condition. Meanwhile, some cases of the AEF phase difference between two stations in the different longitude even in fair weather condition and less ion-aerosol condition existed. In this presentation, we would like to show our preliminary observation results of AEF global variation and Schumann resonances to understand global electric circuit in detail.

Keywords: Global circuit, Atmospheric electric field, ELF electromagnetic waves