

AGE030-07

Room: 202

Time: May 26 10:45-11:00

## The Long-term Diffusion Project at the Grimsel Test Site, Switzerland

Andrew Martin<sup>1\*</sup>, Jussi Ikonen<sup>2</sup>, Marja Siitari-Kauppi<sup>2</sup>, Havlova Vaclava<sup>3</sup>, Akira Hayano<sup>4</sup>

<sup>1</sup>NAGRA, Switzerland, <sup>2</sup>HYRL, Finland, <sup>3</sup>NRI, Czech Republic, <sup>4</sup>JAEA, Japan

The Grimsel Test Site (GTS) ([www.grimsel.com](http://www.grimsel.com)) is an underground rock laboratory located in the crystalline rocks of the Aare Massif in the Swiss Alps. Several projects were recently initiated to study the long-term behavior of radionuclides in a simulated repository near-field and the surrounding host rock involving the setting up of large scale in-situ experiments in the radiation controlled zone of the Grimsel Test Site.

Here an overview is presented of one of such experiment - the Long-Term Diffusion project which is an international partner project\* that aims to verify and understand in-situ processes that control the long-term diffusion of repository relevant radionuclides.

The project is divided into two phases. Phase 1 has been already completed and consisted of four work-packages: (1) an in-situ diffusion experiment in which weakly sorbing and non-sorbing radionuclides (H-3, Na-22, I-131, Cs-134) were circulated and allowed to diffuse into undisturbed rock matrix for almost 800 days; (2) experiments to characterize pore space geometry, including determination of in-situ porosity with C-14 doped PMMA resin for comparison with laboratory derived data; (3) a study of natural tracers to elucidate evidence of long-term diffusion processes; and (4) an investigation of the in-situ matrix diffusion paths for strongly sorbing radionuclides in core material from earlier GTS migration experiments. The present Phase 2 of the project is made up of three work packages: (1) a second monopole involving the circulation of Se-82, Cl-36, H-3 and Ba-133; (2) mock-up tests on block scale granites to derive effective diffusion coefficients for HTO, chloride and selenium; and (3) application of positron-emission-tomography (PET) for the visualization of radionuclide migration through a fracture plane in a Grimsel rock sample.

It is envisaged that by bringing together the results from modeling (predictive and post-mortem), laboratory scale experiments and the in-situ experiments, realistic values for the extent of in-situ matrix diffusion will be determined and further improvements to existing performance assessment codes can be made.

\*JAEA (Japan), University of Helsinki (Finland), NRI (Czech Republic) and NAGRA (Switzerland)

Keywords: Diffusion, Radionuclides, Granite