

Groundwater ages in the southern foot of Mt. Fuji

Yuki Tosaki^{1*}, Norio Tase¹, Kimikazu Sasa², Tsutomu Takahashi³, Yasuo Nagashima²

¹Life & Environ. Sci., Univ. Tsukuba, ²Pure & Appl. Sci., Univ. Tsukuba, ³Appl. Accel. Div., Univ. Tsukuba

Generally, groundwater is highly vulnerable to excess use and contamination, because of its typically long mean residence time. If groundwater depletion and contamination occur, aquifers will require considerably long time for its recovery and purification. Therefore, the knowledge of groundwater flow system and residence time is crucial for the development and sustainable utilization of groundwater resources.

Having an elevation of 3,776 m, Mt. Fuji is one of the largest Quaternary stratovolcano in Japan. It primarily consists of alternating basaltic lava flows and coarse-grained pyroclastic rocks. It can be structurally divided into three parts (i.e. Komitake, Older Fuji and Younger Fuji volcanoes in ascending order). Underlying basement rocks are mainly Neogene marine sediments.

Mt. Fuji holds large groundwater reservoirs in its body because of a total volume of about $2 * 10^9$ m³ of precipitation over a year. Annual precipitation over the Mt. Fuji area ranges from 1500-2000 mm on the northern slope to 2750-3000 mm on the eastern slope. A number of springs distributed over its foot mainly originate from the confined groundwater in permeable part of lava flows which were formed in the early Holocene.

Application of environmental tracers (e.g. CFCs, tritiogenic ³He, SF₆ and ⁸⁵Kr) is one of the most promising approaches for dating groundwater. Bomb-produced ³⁶Cl can also provide a potential dating tool covering the last 60 yrs. In our previous study (Tosaki et al., 2008), we applied ³⁶Cl to estimate residence times of spring waters around Mt. Fuji.

This presentation reports an extension of the previous work, and mainly focuses on the results for the groundwater samples from wells on the southern foot. The ³⁶Cl/Cl ratios in the samples were in the range between $3.3 * 10^{-14}$ and $79.5 * 10^{-14}$, resulting in ³⁶Cl concentrations of $0.5-27.7 * 10^6$ atoms/L. This variation can be ascribed to the differences in groundwater age between aquifers, spanning back to the pre-bomb period.

References:

Tosaki, Y., Tase, N., Sasa, K., Takahashi, T. and Nagashima, Y. (2008): Estimation of groundwater residence time using bomb-produced ³⁶Cl: a case study on spring waters from Mt. Fuji, Japan. In: Proceedings of the XXXVI IAH Congress, pp. 999-1005.