

HQR010-05

会場:展示ホール7別室1

時間: 5月27日11:45-12:00

ネパール、ララ湖堆積物コアを用いた完新世におけるアジアモンスーンの復元

Holocene variability of the Asian monsoon inferred from a sediment core from Lake Rara, western Nepal

中村 淳路^{1*}, 横山祐典², 前杵英明³, 八木浩司⁴, 岡村真⁵, 松岡裕美⁵, 三宅尚⁵, 長田俊樹⁶, 寺村裕史⁶, 山田智輝⁷, Danda Pani Adhikari⁸, Vishnu Dangol⁸, 松崎浩之⁹

Atsunori Nakamura^{1*}, Yusuke Yokoyama², Hideaki Maemoku³, Hiroshi Yagi⁴, Makoto Okamura⁵, Hiromi Matsuoka⁵, Nao Miyake⁵, Toshiki Osada⁶, Hirofumi Teramura⁶, Tomoki Yamada⁷, Danda Pani Adhikari⁸, Vishnu Dangol⁸, Hiroyuki Matsuzaki⁹

¹東大・理・地惑, ²東大 海洋研, ³広島大学 大学院教育学研究科, ⁴山形大学 地域教育学研究科, ⁵高知大学・理, ⁶人間文化研究機構 総合地球環境学研究所, ⁷東北大学大学院文学研究科, ⁸Department of Geology, Tribhuvan Univ., ⁹東大・工

¹EPS, Univ. Tokyo, ²ORI, Univ. Tokyo, ³Hiroshima Univ., ⁴Yamagata Univ., ⁵Kochi Univ.,

⁶National Institutes for the Humanities, ⁷Faculty of Arts and Letters, Tohoku Univ.,

⁸Department of Geology, Tribhuvan Univ., ⁹MALT, Univ. Tokyo

The Asian monsoon is an important component of the Earth's climate system that influences the societal and economic activity of roughly half the world's population (Yancheva et al., 2007). The main driver of large-scale monsoon change over the past 10,000 years has been a slow decrease in summer-time solar radiation (insolation) at 30N owing to changes in Earth's orbit. Even so, monsoon records show abundant evidence of abrupt, stepwise changes on timescales of a century and shorter. The mechanisms behind this behavior are poorly understood. (Overpeck et al., 2007). The inverse correlation between summer and winter monsoons is obvious during the Bolling-Allerod, the Younger Dryas, and the early Holocene. This indicates a link between the monsoon and the movement of ITCZ (Yancheva et al., 2007). However, it is difficult to evaluate the anticorrelation during the late Holocene.

Here we present a new paleoclimatic record from a sediment core recovered in Lake Rara, western Nepal. Lake Rara today lies 3,000m above sea level and has a maximum water depth of 168m. The age model of the sediment core is based on AMS 14C dating of leaves. Concentrations of major elements were measured by X-ray Fluorescence Analysis (XRF). Intensity of chemical weathering in the catchment area was reconstructed by geochemical indexes such as CIA (chemical index of alteration). Bottom-water redox condition was also reconstructed by Mn/Al ratio. Based on the results, we will discuss evolution and variability of the Asian monsoon during the late Holocene.

Acknowledgement

This study was supported by the research project titled Environmental Change and the Indus Civilization (project 3-3) managed by Research Institute for Humanity and Nature, Japan.

Yancheva, G. et al., Nature 445, 74-77 (2007)

Overpeck, J.T. et al., Nature 445, 270-271 (2007)

キーワード: モンスーン, ララ湖, ネパール, 完新世, XRF

Keywords: monsoon, Lake Rara, Nepal, Holocene, XRF