

## 高速太陽風通過時における放射線帯加速

### Acceleration of outer belt electrons associated with high speed streams

三好 由純<sup>1\*</sup>, 片岡 龍峰<sup>2</sup>

Yoshizumi Miyoshi<sup>1\*</sup>, Ryuho Kataoka<sup>2</sup>

<sup>1</sup>名古屋大学太陽地球環境研究所, <sup>2</sup>東京工業大学

<sup>1</sup>STEL, Nagoya University, <sup>2</sup>Tokyo Institute of Technology

Flux enhancements of relativistic electrons in the outer belt tend to occur associated with high-speed streams, and a large flux enhancement is observed during the predominantly southward IMF. Our statistical studies revealed that 90%(50%) of the fast coronal hole streams display a large flux enhancement at GEO when the southward (northward) IMF is dominant. From the standpoint of the internal acceleration by whistler mode chorus waves, this IMF dependence can be understood as follows. The internal acceleration by wave-particle interactions is especially effective when a continuous source of hot electrons can be maintained to produce chorus waves for the several day periods. Continuous hot electron injections are enhanced during a prolonged period of intense convection/substorms, which are driven by the southward IMF in high-speed streams. The statistical studies show clear differences of hot electrons, whistler mode chorus waves and substorm/convection between the southward and northward dominant IMF in high-speed streams, which are consistent with the internal acceleration process.

キーワード:放射線帯,粒子加速,高速コロナホール流

Keywords: radiation belts, particle acceleration, high speed streams