

PEM021-P08

会場:コンベンションホール

時間: 5月24日17:15-18:45

GEMSIS-電離圏:擾乱時のグローバルな電離圏電場ポテンシャル分布の導出へ向けて

GEMSIS-Ionosphere: Toward the understanding of global distribution of the ionospheric potential during disturbed time

海老原 祐輔^{1*}, 菊池 崇², 家田 章正², 新堀 淳樹², 平木 康隆², 堀 智昭², 西谷 望², 樋口 知之³, 上野 玄太³, Ohtani Shinichi⁴

Yusuke Ebihara^{1*}, Takashi Kikuchi², Akimasa Ieda², Atsuki Shinbori², Yasutaka Hiraki², Tomoaki Hori², Nozomu Nishitani², Tomoyuki Higuchi³, Genta Ueno³, Shinichi Ohtani⁴

¹名古屋大学 高等研究院, ²名古屋大学 太陽地球環境研究所, ³統計数理研究所, ⁴ジョンズホプキンス大学応用物理研究所

¹IAR, Nagoya University, ²STEL, Nagoya University, ³ISM, ⁴JHU/APL

In phase 1 of the GEMSIS (Geospace Environment Modeling System for Integrated Studies)-Ionosphere project, the primary goal was set to reconstruct the global distribution of the ionospheric electric potential for the purpose of understanding the Sun-Earth system. Because of irregularly distributed observatories and insufficient number of them, it has been a challenging work to reconstruct the electric potential distribution. We initiated two different challenges. One is an inductive scheme using data from the SuperDARN radars and ground magnetic fields. By applying a new base model, a global-scale distribution of the electric potential will be obtained for the storm-time in which the convection cell is largely expanded to lower latitudes. The other challenge is a deductive scheme by applying Ohm's law for given electric currents flowing along a field line (field-aligned current) and the ionospheric conductance. One of the advantages of our deductive scheme is to solve the potential in the whole sphere under the realistic geomagnetic field. We developed a realistic model of the large-scale field-aligned current on the basis of 190,000 data sets of field-aligned current observations obtained by the DMSP and DE2 satellites. The data were sorted by the solar wind condition and tilt angle, providing realistic field-aligned currents having sharp boundaries. Data analysis efforts have also been made in terms of quantitative evaluation of transmission of electromagnetic energy, auroral breakup and magnetotail reconnection during substorms, inner magnetosphere-ionosphere coupling, and magnetosphere-ionosphere coupling system.

キーワード:電離圏電場ポテンシャル,磁気嵐,サブストーム,沿磁力線電流,モデリング

Keywords: Ionospheric electric potential, Magnetic storms, Substorms, Field-aligned current, Modeling