

## Effect of the solar-wind proton entry into the deepest lunar wake

Masaki N Nishino<sup>1\*</sup>, Masaki Fujimoto<sup>1</sup>, Yoshifumi Saito<sup>1</sup>, Shoichiro Yokota<sup>1</sup>,  
Yoshiya Kasahara<sup>2</sup>, Yoshiharu Omura<sup>3</sup>, Yoshitaka Goto<sup>2</sup>, Kozo Hashimoto<sup>3</sup>,  
Atsushi Kumamoto<sup>4</sup>, Takayuki Ono<sup>4</sup>, Hideo Tsunakawa<sup>5</sup>, Masaki Matsushima<sup>5</sup>,  
Futoshi Takahashi<sup>5</sup>, Hidetoshi Shibuya<sup>6</sup>, Hisayoshi Shimizu<sup>7</sup>, Toshio Terasawa<sup>8</sup>

<sup>1</sup>ISAS/JAXA, <sup>2</sup>Kanazawa University, <sup>3</sup>RISH, Kyoto University, <sup>4</sup>Tohoku University,

<sup>5</sup>Tokyo Institute of Technology, <sup>6</sup>Kumamoto University, <sup>7</sup>ERI, University of Tokyo, <sup>8</sup>ICRR, University of Tokyo

We study effect of the solar wind (SW) proton entry deep into the near-Moon wake that was recently discovered by the SELENE mission. It has been accepted that the high-speed electrons in the ambient SW determine the lunar night-side environment. Therefore, previous lunar-wake models are based on the electron domination, and no effect of SW proton entry on the near-Moon wake environment has been taken into account so far. Recent SELENE observations revealed that a part of the SW protons are reflected at the lunar dayside surface and picked-up by the SW electric field (Saito et al., GRL, 2008), and some of them access the deepest lunar wake; this process is called type-II entry (Nishino et al., GRL, 2009). Here we show that the type-II entry of SW protons forms proton-governed region (PGR) to drastically change the electromagnetic environment of the lunar wake. Broadband electrostatic noise found in the PGR is manifestation of electron two-stream instability, which is attributed to the counter-streaming electrons absorbed from the ambient SW to maintain the quasi-neutrality. An acceleration of the absorbed electrons up to about 1 keV means a superabundance of positive charges of  $10^{-4}$ - $10^{-7}$  cm<sup>-3</sup> in the near-Moon wake, which should be immediately canceled out by the incoming high-speed electrons. This is a general phenomenon in the lunar wake, because PGR does not necessarily require peculiar SW condition for its formation.

Keywords: Lunar wake, Solar wind plasma, Plasma waves, Particle acceleration