

PEM025-P26

会場:コンベンションホール

時間: 5月26日17:15-18:45

ISS-IMAP搭載VISIによる大気光観測：開発の現状と観測シミュレーション(3)

Visible airglow observation by VISI on ISS-IMAP: Current status of development and simulation of airglow measurement (3)

坂野井 健^{1*}, 秋谷 祐亮², 大塚 雄一³, 山崎 敦⁵, 阿部 琢美⁵, 齊藤 昭則², 鈴木 睦⁵,
武山 芸英⁴, 小淵 保幸⁴, 田口 真⁶, 江尻 省⁷, 久保田 実⁸, 吉川 一朗⁹, 山本 衛¹⁰

Takeshi Sakanoi^{1*}, Yusuke Akiya², Yuichi Otsuka³, Atsushi Yamazaki⁵, Takumi Abe⁵,
Akinori Saito², Makoto Suzuki⁵, Norihide Takeyama⁴, Yasuyuki Obuchi⁴, Makoto Taguchi⁶,
Mitsumu Ejiri⁷, Minoru Kubota⁸, Ichiro Yoshikawa⁹, Mamoru Yamamoto¹⁰

¹東北大学大学院理学研究科, ²京都大学大学院理学研究科, ³名古屋大学太陽地球環境研究所, ⁴(株)ジェネシア,
⁵宇宙航空研究開発機構宇宙科学研究本部, ⁶立教大学理学部, ⁷国立極地研究所, ⁸情報通信研究機構,
⁹東京大学大学院理学研究科, ¹⁰京都大学生存圏研究所

¹Grad Sch. of Science, Tohoku University, ²Grad Sch. of Science, Kyoto University, ³STEL, Nagoya University,
⁴Genesis Corp., ⁵JAXA/ISAS, ⁶Faculty of Science, Rikkyo University, ⁷NIPR, ⁸NICT,
⁹Sch. of Science, University of Tokyo, ¹⁰RISH, Kyoto University

The ISS-IMAP mission is one of the constituents of the Japanese Experiment Module (JEM) 2nd stage plan which will be launched in 2011 or 2012 onto the International Space Station (ISS). The main scientific subject of this mission is to clarify the energy and physical transfer processes in the boundary region between earth's atmosphere and space with the visible spectrometer and extra ultraviolet imagers.

We have been developing the visible imaging spectrometer instrument (VISI) on ISS-IMAP. VISI will measure three nightglow emissions; O (630 nm, altitude 250 km), OH Meinel band (730 nm, altitude 87km), and O₂ (0-0) atmospheric band (762 nm, altitude 95 km). We designed a bright (F/0.9), wide-angle (field-of-view 90 degrees) objective lens. To subtract background contaminations from clouds and ground structures, VISI will perform a stereoscopic measurement by taking two line-scans of the airglow spectra in 45 deg. forward and 45 deg. backward directions. Each field-of-view is faced perpendicular to the orbital plane, and its width is about 550 km mapping to an altitude of 100 km. The phase velocity of airglow wave structure is also estimated from the difference between forward and backward data. We will obtain a continuous line-scanning image for all emissions line from + 51 deg to -51 deg. in geographic latitude by the successive exposure cycle with a time interval of 1 - several sec. From the simulation of airglow observation with a simple sine-curve airglow pattern including noises, we found that the wave structure in airglow can be identified for the three emissions. In this presentation, the current status of instrumental development, system design, operation plan, and simulation of airglow observations will be reported.

キーワード:国際宇宙ステーション,きぼう,熱圏,電離圏,開発,光学

Keywords: ISS, JEM, thermosphere, ionosphere, development, optics