

PEM027-06

会場:ファンクションルームB

時間: 5月24日15:00-15:15

あけぼの衛星を用いた太陽周期変動型電子放射線帯モデル

Electron Radiation Belt Model Varying Through The Solar Cycle Using Akebono Satellite

林 幹子¹, 中村 雅夫^{1*}, 浅井 佳子², 高田 拓³, 三好 由純⁴, 高島 健³, 松岡 彩子³,
長井 嗣信²

Mikiko Hayashi¹, Masao Nakamura^{1*}, Keiko T. Asai², Taku Takada³, Yoshizumi Miyoshi⁴,
Takeshi Takashima³, Ayako Matsuoka³, Tsugunobu Nagai²

¹大阪府立大学, ²東京工業大学, ³宇宙航空研究開発機構, ⁴名古屋大学太陽地球環境研究所

¹Osaka Prefecture University, ²Tokyo Institute of Technology, ³Japan Aerospace Exploration Agency,
⁴STEL, Nagoya University

The radiation belts are regions of high-energy particles, mainly protons and electrons, trapped by the Earth's magnetic field. The radiation belt models are important for the design and operation of spacecrafts because the high-energy particles can damage the satellites, e.g., directly by single event effects and on prolonged exposure, and can be a hazard to astronauts. The annual averaged high-energy electron flux of the radiation belt varies with solar activities which undergo 11-year solar cycles. We have examined the variations of the electron radiation belts with the solar and magnetospheric activities using data from the Radiation Monitor (RDM) of the Akebono satellite. The Akebono satellite has been in the highly elliptical low Earth orbit over the last two decades and measured the almost whole region of the inner belt and the high latitude region of the outer belt. We have proposed the new electron radiation belt model varying through the solar cycle based on the unique measurement of the Akebono satellite, which has been continuously done for almost two solar cycles.

キーワード:電子放射線帯モデル,あけぼの衛星,太陽周期

Keywords: Electron radiation belt model, Akebono satellite, Solar cycle