

PEM028-03

会場:展示ホール7別室1

時間: 5月23日14:33-14:50

超高強度レーザーによる宇宙線加速模擬実験

Model Experiment of Cosmic Ray Acceleration due to an Incoherent Wakefield Induced by an Intensive Laser Pulse

藏満 康浩^{1*}, 中新信彦¹, 近藤公伯², 坂和洋一¹, 森芳孝³, 三浦永祐⁵, 谷本壯¹, 中村浩隆¹, 反保元信², 児玉了祐¹, 北川米喜³, 三間闇興¹, 田中和夫¹, 星野真弘⁴, 高部英明¹

Yasuhiro Kuramitsu^{1*}, Nobuhiko Nakanii¹, Kiminori Kondo², Youichi Sakawa¹, Yoshitaka Mori³, Eisuke Miura⁵, Tsuyoshi Tanimoto¹, Hirotaka Nakamura¹, Motonobu Tanpo², Ryosuke Kodama¹, Yoneyoshi Kitagawa³, Kunioki Mima¹, Kazuo Tanaka¹, Masahiro Hoshino⁴, Hideaki Takabe¹

¹大阪大学, ²日本原子力研究開発機構, ³光産業創成大学院大学, ⁴産業技術総合研究所, ⁵東京大学

¹Osaka University, ²Japan Atomic Energy Agency, ³Graduate School for the Creation of New, ⁴National Institute of Advanced Industria, ⁵University of Tokyo

We performed a model experiment of cosmic ray acceleration by using intensive laser pulses. Large amplitude light waves are considered to be excited in the upstream regions of relativistic astrophysical shocks and the wakefield acceleration of cosmic rays can take place. We model such shock environments in a laboratory plasma by substituting an intensive laser pulse for the large amplitude light waves. A plasma tube, which is created by imploding a hollow polystyrene cylinder, was irradiated by an intensive laser pulse. Nonthermal electrons were generated by the wakefield acceleration and the energy distribution functions of the electrons have a power law component with an index of ~ 2.5 . The maximum attainable energy of the electrons in the experiment is discussed by a simple analytic model. In the incoherent wakefield the maximum energy can be much larger than in the coherent field due to the momentum space diffusion or the energy diffusion of electrons.

キーワード:宇宙線,粒子加速,衝撃波,実験室宇宙物理

Keywords: cosmic rays, acceleration of particles, shock waves, laboratory astrophysics