

ひのでXRTによるコロナホールの観測

Structure and Evolution of Coronal Holes Observed with Hinode XRT

坂尾 太郎^{1*}, 成影典之², 鹿野良平²

Taro Sakao^{1*}, Noriyuki Narukage², Ryouhei Kano²

¹宇宙航空研究開発機構・宇宙科学研究本部, ²自然科学研究機構・国立天文台

¹ISAS/JAXA, ²NAOJ

We report structure and its dynamic evolution, both in intensity and temperature, of equatorial coronal holes observed with the X-Ray Telescope (XRT) aboard Hinode. Coronal hole that has been studied primarily is the one located near the disk center on 27-29 April 2008, when no active region was present on the solar disk.

This coronal hole was observed extensively, with a pair of Al/Mesh and Titanium-on-Polyimide (Ti/Poly) filters with each filter pair taken every 90 seconds. As this coronal hole was located near the disk center, we expect that, along the line of sight, X-ray emission from the hole was less contaminated by that from (diffuse) X-ray loops from the surrounding quiet Sun as compared to coronal holes near the limb, hence more suited to study physical conditions inside coronal holes.

When seen with the Al/Mesh filter, the coronal hole is by no means very dark (i.e., less contrast to the quiet Sun than the one seen with thicker filters), but instead exhibits frequent transient brightenings across the hole with time scale of a few minutes. The observed brightenings indicate frequent emergence and submergence of magnetic flux near the base of the coronal hole, from the lower atmosphere. The boundary portions of the coronal hole are generally low in temperature (at, or slightly lower than, 1 MK) with the lowest-temperature portion (< ~0.8 MK) created in association with apparent change in magnetic configuration near the boundary of the coronal hole.

Details of evolution in temperature and intensity structure of coronal holes will be discussed.

キーワード:太陽,ひので,コロナホール, X線

Keywords: Sun, Hinode, Coronal holes, X-rays