Japan Geoscience Union Meeting 2010

(May 23-28 2010 at Makuhari, Chiba, Japan)

©2009. Japan Geoscience Union. All Rights Reserved.



PEM032-P01

会場:コンベンションホール

時間: 5月25日17:15-18:45

衝撃波静止系による準垂直衝撃波の2次元粒子シミュレーション

Two-dimensional particle simulation of a quasi-perpendicular shock with a shock rest frame model

木谷 佳隆^{1*}, 梅田 隆行¹, 山尾 政博¹, 山崎 了², 荻野 竜樹¹

Yoshitaka Kidani^{1*}, Takayuki Umeda¹, Masahiro Yamao¹, Yamazaki Ryo², Tatsuki Ogino¹

¹名古屋大学太陽地球環境研究所, ²広島大学大学院理学研究科物理専攻

¹STEL, Nagoya Univ., ²Hiroshima Univ.

A full particle simulation study is carried out on the reformation of a quasi-perpendicular collisionless shock with a relatively low Alfven Mach number (MA = 5). Recent self-consistent hybrid and full particle simulations have demonstrated that ion kinetics are essential for the non-stationarity of perpendicular collisionless shocks, which means that physical processes due to ion kinetics modify the shock jump condition for fluid plasmas. However, it is a heavy task to conduct large-scale full particle simulations of collisionless shocks to include electron kinetics. In the present study, we have performed a two-dimensional (2D) electromagnetic full particle simulation with a "shock-rest-frame model". The simulation domain is taken to be larger than the ion inertial length in order to include full kinetics of both electrons and ions. We discuss the effect of electron kinetics to the shock reformation at a quasi-perpendicular shock.

Keywords: shock wave, fulll-particle simulation, quasi-perpendicular, low Alfven Mach number