

Japan Geoscience Union Meeting 2010

(May 23-28 2010 at Makuhari, Chiba, Japan)

©2009. Japan Geoscience Union. All Rights Reserved.



PEM033-05

会場: 303

時間: 5月27日11:57-12:14

超新星爆発への太陽地球環境応答

Response of the solar-terrestrial environment to supernova explosions

片岡 龍峰^{1*}

Ryuho Kataoka^{1*}

¹東工大

¹Tokyo Tech

Nearby supernova explosions at 10 pc distance from the Sun put the Earth inside the supernova remnant (SNR), and the severe condition can be a possible origin of Snowball Earth and the birth of complex life. Highly compressed and turbulent heliosphere is predicted from 3D magnetohydrodynamic (MHD) simulation of the nearby SNR collision to the Sun-Earth system. Richtmyer-Meshkov instability at nose may provide a local source of the turbulence associated with the Kelvin-Helmholtz instability in the inner and outer heliosheaths. Asymmetric shape of the heliosphere is expected due to the Parker spiral of interplanetary magnetic field. As a result, it is shown that roughly thousand times larger cosmic ray flux than the present value is predicted to continue for a few thousand years associated with the nearby supernova explosions at 10 pc, while the cosmic ray flux would be limited only a few times larger than the present level for the 100 pc distance cases.