

## 磁気プラズマセイルにおける人工磁気圏展開と太陽風相互作用

### Solar wind interaction and magnetic inflation for small scale artificial magnetosphere around a magneto-plasma sail

森高 外征雄<sup>1\*</sup>, 白井 英之<sup>1</sup>, 沼波 政倫<sup>2</sup>, 梶村 好宏<sup>3</sup>, 篠原 育<sup>4</sup>, 中村 雅夫<sup>5</sup>

Toseo Moritaka<sup>1\*</sup>, Hideyuki Usui<sup>1</sup>, Masanori Nunami<sup>2</sup>, Yoshihiro Kajimura<sup>3</sup>, Iku Shinohara<sup>4</sup>, Masao Nakamura<sup>5</sup>

<sup>1</sup>神戸大学大学院工学研究科, <sup>2</sup>核融合科学研究所, <sup>3</sup>京都大学 生存圏研究所,  
<sup>4</sup>宇宙科学研究本部 宇宙科学情報解析研究系, <sup>5</sup>大阪府立大学大学院工学研究科

<sup>1</sup>Kobe University, <sup>2</sup>National Institute for Fusion Science, <sup>3</sup>Kyoto University, <sup>4</sup>Japan Aerospace Exploration Agency, <sup>5</sup>Osaka Prefecture University

Magneto-plasma sail is one of the next generation flight systems to the deep space[1,2]. The propulsive force is obtained from interaction between the solar wind and the artificial magnetosphere created around the spacecraft. Previous hybrid and Hall simulations have suggested that a magnetosphere with a spatial scale comparable to the ion inertia length is necessary for effective interaction with the solar wind. On the other hand, realistic size coil of a few meters in radius is expected to create only an electron scale magnetosphere of a few kilo meters large or less. To solve this problem, magnetic inflation via plasma injection from the spacecraft has been proposed[3]. However the interactions of the electron-scale magnetosphere with the injection plasma and the solar wind are still poorly understood.

Two dimensional full particle-in-cell simulations have been performed to investigate the magnetic inflation and the solar wind interactions for the small scale artificial magnetospheres. The magnetic inflation processes is characterized by the gyration motion of the injected ions. Ion rich regions are formed at the turning around position of the injected ions and electrons flow into this position carrying the field lines of the original magnetosphere. As a result, the magnetosphere is deformed effectively when the spatial size of the original magnetosphere is comparable to the ion gyration radius. For the electron scale magnetosphere, the inflation process is confined to the smaller region before the ion gyration point as a result of the interaction between background electrons and the injected ions.

The solar wind interactions and the resulting gain of the propulsive force for the electron scale and the deformed magnetospheres will also be discussed by comparison to that for the ion scale magnetosphere.

[1] R. M. Zubrin and R. G. Andrews, *Journal of Spacecraft and Rockets*, 28, 197 (1991)

[2] H. Yamakawa, I. Funaki, Y. Nakayama et al., *Acta Astronautica* 59, 777, (2006).

[3] R. M. Winglee et al., *Journal of Geophysical Research* 105, 67 (2000).

キーワード: PICシミュレーション, 磁気プラズマセイル, 小型磁気圏, 磁気インフレーション

Keywords: PIC simulation, magneto-plasma sail, small-scale magnetosphere, magnetic inflation