

PPS001-09

会場: 304

時間: 5月23日11:23-11:36

火星における中小規模擾乱のダスト巻き上げにおける影響:大気大循環モデル実験

Effects of small and medium scale disturbances on the dust lifting on Mars: general circulation model experiments

高橋 芳幸^{1*}, 林祥介², 小高正嗣³

Yoshiyuki O. Takahashi^{1*}, Yoshi-Yuki Hayashi², Masatsugu Odaka³

¹惑星科学研究センター, ²神戸大学大学院理学研究科, ³北海道大学大学院理学研究院

¹Center for Planetary Science, ²Kobe University, ³Hokkaido University

High resolution experiments of Mars atmospheric circulation by using a general circulation model have been conducted to address the features of small and medium scale disturbances and its effects on dust lifting. The mechanisms of dust lifting on Mars have not been understood well, while dust suspended in the atmosphere is one of the most important factors that characterize the thermal structure and circulation of the Martian atmosphere. The high resolution simulations that can resolve the small and medium scale disturbances over the globe may provide important insights into the lifting mechanism of dust on Mars. In this study, the experiments with several resolutions are conducted and the effects of small and medium scale disturbances are examined by examining the frequency distribution of surface stress in the model.

The model used in this study consists of the dynamical core of AFES, and the physical processes introduced from the Mars GCM which has been developed by our group so far. AFES is based on CCSR/NIES AGCM 5.4.02, and is optimized to conduct high resolution experiments on the Earth Simulator. The introduced physical processes include the radiative, the turbulent mixing, and the surface processes. In addition, the dust lifting process is implemented to diagnose the dust lifting rate in the model. The dust lifting process is the same as one of "threshold-sensitive surface stress lifting" parameterizations proposed by Newman et al. (2002). This parameterization is a GCM implementation of the process of dust lifting by the surface wind, whose characteristics is that dust is not lifted unless the surface friction velocity exceeds a certain threshold value. By the use of this GCM, we performed simulations at northern fall season with resolutions of T79L96, and T159L96, which are equivalent to about 89 and 44 km grid size, respectively, and with 96 vertical layers. In these experiments, the dust is treated as a radiative passive material for the ease of interpretation of the results.

The examination of frequency distribution of surface stress in the results of T79L96 and T159L96 experiments show that the high surface stress tail of distribution is found in T159L96 experiment compared to that in T79L96 experiment. Clearly, this is an appearance of effects of small and mediums scale disturbances that are resolved in the T159L96 experiment and are not resolved in the T96L96 experiment. In the presentation, the effects of topography will be discussed by using the results of experiment with no topography.

キーワード:火星,惑星大気,大気大循環モデル,中小規模擾乱,地球シミュレータ

Keywords: Mars, planetary atmosphere, general circulation model,

small and medium scale disturbance, Earth simulator