

アモルファス氷への真空紫外光照射による水分子脱離メカニズム

A desorption mechanism of water following vacuum-ultraviolet irradiation of amorphous solid water

羽馬 哲也^{1*}, 横山 正明¹, 藪下 彰啓¹, 川崎 昌博¹, 渡部 直樹²

Tetsuya Hama^{1*}, Yokoyama Masaaki¹, Akihiro Yabushita¹, Masahiro Kawasaki¹, Naoki Watanabe²

¹京都大学大学院工学研究科分子工学専攻, ²北海道大学低温科学研究所

¹Kyoto University, ²Hokkaido University




Figure 1 Schematic illustration for photodesorption of H₂O molecule from amorphous solid water via “kick-out” mechanism by a hot H atom released from photodissociation of amorphous water ice. The numbers dedicate for sequence of the reaction.

Water ice is the major solid component in a variety of astrophysical environments, ranging from cold and dense molecular clouds, the small solar system objects such as Kuiper Belt objects, comets, to icy satellites in the outer solar system. In such cold regions where the thermal evaporation of ice is negligible, photodesorption plays a dominant role in consuming ice. Several experimental studies of the photodesorption of H₂O molecules from ice surfaces have been performed. However, these studies did not reveal the desorption mechanism or the energy partitioning of photodesorbed H₂O molecules, which is important for detailed modeling of gas-ice surface interactions, and/or as possible observational signatures of photodesorption.

In this study, photodesorption of H₂O molecules from amorphous solid water at 90 K has been investigated, in order to probe the photodesorption mechanism. Using resonance-enhanced multiphoton ionization technique, we have measured the translational and rotational energy distributions of photodesorbed H₂O molecules following vacuum-ultraviolet (VUV) laser irradiation of amorphous solid water.

The measured translational and rotational temperatures are 1800 K and 300 K, respectively. These energies are in good accord with those predicted by classical molecular dynamics calculations for H₂O photodesorption by a kick-out mechanism following absorption of a single UV photon, that is, an H₂O molecule is ejected from the ice by momentum transfer, mainly to the oxygen atom in H₂O, from an hot H atom released by photodissociation of a neighboring H₂O, i.e., reactions (1) and (2);

Photodissociation: H₂O(ice) + VUV → hot H + OH, dH = 541.6 kJ mol⁻¹(1)

Water desorption: hot H + H₂O(ice) → H + H₂O(gas). dH = 44.5 kJ mol⁻¹(2)

キーワード:アモルファス氷,光脱離,水分子,彗星,カイパーベルト天体群

Keywords: amorphous solid water, photodesorption, water molecule, comet, Kuiper Belt objects