

PPS004-08

会場: 201A

時間: 5月25日11:00-11:15

太陽系外の逆行惑星の形成

Formation of retrograde extrasolar planets

長沢 真樹子^{1*}, 井田 茂¹

Makiko Nagasawa^{1*}, Shigeru Ida¹

¹東京工業大学

¹Tokyo Institute of Technology

About 25% of the discovered extrasolar planets have orbital periods smaller than 1 week. These are called as the close-in planets. Jovian or Neptunian type planets in this location are likely to be formed at large distance from their host stars and migrated to shorter period orbits. Sky projected angles between a stellar equator and a planetary orbit are measured for 20 transiting planet using Rossiter-McLaughlin technique. About half of them are in highly inclined orbits. This misalignment is hardly explained by Type-II migration, which is most contributing mechanism to planetary migration.

Starting with three planets orbiting around a solar mass star in circular orbits, I have numerically investigated how planetary orbits evolve by a combination effect of mutual scattering and tidal circularization for understanding the inclination distribution. Dynamical tides corresponding to planetary normal modes and inertial modes, relativity, gravity of the central star, and mutual gravity of planets are included in simulations.

I obtain the close-in planets in about 24% of runs. The formed close-in planets have a broad range of orbital inclinations. The probability of retrograde orbit is 29%. Approximately 83% of the close-in planet are formed during planet-planet scattering phase. That means they tend to have very short tidal evolution timescale and they tend to be totally circularized. On the other hand, about 17% of the close-in planets are formed after the scattering phase. They tend to have longer tidal evolution timescale. Since their eccentricities are gradually increased by the secular interaction or the Kozai mechanism exchanging the inclination and eccentricity, they tend to be in prograde orbits or in near polar retrograde orbits. The retrograde planet with moderate inclination and high eccentricity is rare, since they need to be scattered to very limited region of orbital parameters during orbital instability stage.

This work was supported by MEXT-KAKENHI(21740324) 'Grant-in-Aid for Young Scientists (B)' and MEXT's program 'Promotion of Environmental Improvement for Independence of Young Researchers' under the Special Coordination Funds for Promoting Science and Technology.

キーワード:系外惑星,軌道進化

Keywords: extrasolar planet, orbital evolution