## **Japan Geoscience Union Meeting 2010**

(May 23-28 2010 at Makuhari, Chiba, Japan)

©2009. Japan Geoscience Union. All Rights Reserved.



PPS010-04

会場: 201A

時間: 5月27日14:30-14:45

## 周惑星円盤へのガス降着流と衛星系形成

## Accretion Flow onto Circum-Planetary Disks and Satellite Formation

谷川 享行1\*, 大槻 圭史2, 町田 正博3, 小林 浩4

Takayuki Tanigawa<sup>1\*</sup>, Keiji Ohtsuki<sup>2</sup>, Masahiro Machida<sup>3</sup>, Hiroshi Kobayashi<sup>4</sup>

1北大低温研, 2神戸大・理, 3国立天文台, 4イエナ大学

<sup>1</sup>ILTS, Hokkaido Univ., <sup>2</sup>Graduate School of Science, Kobe Univ., <sup>3</sup>NAOJ, <sup>4</sup>Jena Univ.

Most of major natural satellites around the giant planets in our solar system are classified into "regular satellites", whose orbits are nearly in circular orbits and in the equatorial plane of the parent bodies. The satellites are thus thought to form in circum-planetary disks. However, the origin and the properties of circum-planetary disks are not well understood. Recently, some hydrodynamic simulations showed that gas flow accreting toward the planet forms disk-like structure in the course of gas capturing process to be gas giant planets, thus circum-planetary disks are inevitably formed as by-products of the formation of gas-giant planets.

In order to form satellites, however, solid material needs to be supplied to the circum-planetary disks. The way of supplying solid material can be divided into two. For small objects, they move with gas because of strong gas drag. For large objects, they are basically decoupled with gas motion and affected weakly by gas-drag. In the period of giant planet formation, collision between relatively large objects are frequently occurred and large fraction of solid may be in the form of small dust particles, which couple well with gas motion.

In this study, we conduct high-resolution gas accretion flow from proto-planetary disks to gas giant planets in order to understand structure of circum-planetary disks and resultant dust accretion in the context of satellite formation. We analyze the accretion flow in detail and show that (1) gas near the mid plane in proto-planetary disks are difficult for accretion to the planets, and as a result, sedimentary dust are difficult to be supplied to circum-planetary disks, (2) gas that is going to accrete to the planets is fallen directly within less than 0.05 Hill radius from the planets, meaning that disk-like structure outside of the radius cannot accrete to the planet and dust in the region would not be material to form satellites.

キーワード:衛星,巨大惑星,円盤,流体計算

Keywords: satellites, giant planets, disks, hydrodynamic simulation