

U004-02

会場: 302

時間: 5月24日09:20-09:40

東海下の構造と地震発生に果たす水の役割

Effects of fluids dehydrated from subducting oceanic crust on earthquakes and slow slips in Tokai, Japan

加藤 愛太郎^{1*}, 飯高 隆¹, 生田 領野², 吉田 康宏³, 勝俣 啓⁴, 岩崎 貴哉¹, 酒井 慎一¹,
山岡 耕春⁵, 渡辺 俊樹⁵, 國友孝洋², 山崎 文人⁵, 津村 紀子⁶, 野崎 謙治⁶, 高橋 福助⁶,
大久保 慎人⁷, 鈴木 貞臣⁷, 平田 直¹

Aitaro Kato^{1*}, Takashi Iidaka¹, Ryoya Ikuta², Yasuhiro Yoshida³, Kei Katsumata⁴,
Takaya Iwasaki¹, Shin'ichi Sakai¹, Koshun Yamaoka⁵, Toshiki Watanabe⁵,
Takahiro Kunitomo², Fumihito Yamazaki⁵, Noriko Tsumura⁶, Kenji Nozaki⁶,
fukusuke takahashi⁶, Makoto OKUBO⁷, Sadaomi Suzuki⁷, Naoshi Hirata¹

¹東京大学地震研究所, ²静大, ³気象研, ⁴北大, ⁵名大, ⁶千葉大, ⁷東濃地震科研

¹ERI, University of Tokyo, ²Shizuoka Univ., ³MRI, ⁴Hokkaido Univ., ⁵Nagoya Univ., ⁶Chiba Univ., ⁷TRIES

Episodic slow slips and non-volcanic tremors have been widely detected near the down-dip limit of megathrust earthquakes around the Circumstance Pacific subduction zones. Although the ETS is generally supposed to be attributed to high pressurized fluids, the geological origin of the high fluid pressures in the ETS area has not been figured out in terms of the dehydration metamorphic reactions of the oceanic crust. Here, we show that the subducting oceanic crust of the Philippine Sea plate is undergoing progressive metamorphic dehydration-reactions, based on depth changes of high-resolution seismic images. The Poisson-ratio and the degree of the low-velocity anomaly within the oceanic crust gradually increases from the shallow depths of the down-dip limit, and reaches the peak at centroid of the moment-released area by the long-term slow slip. The Poisson-ratio maintains high value showing a patchy structure at depths greater than the slow slip area, where the non-volcanic low-frequency earthquakes align along the plate interface. They are located beneath a partially serpentized-mantle wedge corner aligning along the plate interface. The oceanic crust then shows rapid reduction of the Poisson-ratio at the greater depths accompanying with abundant intraslab seismicity, where the oceanic basalt could transform to the amphibole-elcogite phase. The slow slip was induced by the near lithostatic fluid pressures generated by an impermeable body underplated beneath the overlying plate. In contrast, the non-volcanic LFEs could be a shear slip of the mantle peridotite contact plane triggered by the locally high-fluid pressures.