

U004-P24

会場:コンベンションホール

時間: 5月24日17:15-18:45

アメリカ, グリーンノブの岩水マントルかんらん岩捕獲岩の地球化学的特徴

Geochemical features of hydrated mantle peridotite xenoliths from the Green Knobs diatreme, USA

石丸 聡子^{1*}, 荒井 章司¹, 石田義人¹, ダグラス スミス²

Satoko Ishimaru^{1*}, Shoji Arai¹, Yoshito Ishida¹, Douglas Smith²

¹金沢大学理工研究域自然システム学系, ²テキサス大学地球学教室

¹Dept. Earth Sci., Kanazawa Univ., ²Dept. Earth Sci., Texas Univ. Austin

Mantle peridotites beneath the Colorado Plateau have been well documented by several workers (e.g. McGetchin and Silver, 1972; Smith and Levy, 1976), and are mostly spinel-garnet lherzolite to harzburgite with minor garnet peridotites. They are characterized by the presence of hydrous minerals, i.e. amphiboles (pargasite to tremolite and sodic tremolite), chlorite, titanoclinohumite and antigorite. There has been a controversy about the source of fluids responsible for the hydration. We determined trace-element characteristics of all constituent minerals of hydrated peridotite xenoliths from the Green Knob diatreme (kimberlitic breccia) to constrain the origin of the fluids.

We examined selected 8 ultramafic xenoliths (3 lherzolites, 3 harzburgites, 1 websterite, and 1 orthopyroxenite), which contain various hydrous minerals (amphiboles, chlorite and titanoclinohumite). The main rock types of the xenolith from Green Knobs are granite, metamorphic rocks, and ultramafics, only 1 % in volume of all (Smith and Levy, 1976). These xenoliths contain sulfide globules, and ilmenite grains are also rarely observed. The chondrite-normalized REE patterns of clinopyroxene slope gently from HREE (Lu) to MREE (Sm) and are high in LREE. Some clinopyroxenes are enriched in Sr and other LILE but depleted in HFSE, i.e. Zr and Ti. These features are inherited from the metasomatic agent. The trace-element abundances of all clinopyroxenes are almost the same except one lherzolite sample that contains garnet pseudomorphs.

We try to discuss the metasomatic event, especially in hydration process, beneath Green Knobs from the geochemical features.

キーワード:かんらん岩捕獲岩, マントルウェッジ, 加水作用, マントル交代作用

Keywords: peridotite xenoliths, mantle wedge, hydration, mantle metasomatism