

BAO001-08

会場: 301B

時間: 5月24日11:00-11:15

大気圏上空微生物に基づく生命の宇宙伝播仮説の評価

## Upper Atmospheric Microorganisms Based Evaluation of Panspermia Hypothesis

Yang Yinjie<sup>1\*</sup>, 横堀 伸一<sup>1</sup>, 山岸 明彦<sup>1</sup>

Yinjie Yang<sup>1\*</sup>, Shin-ichi Yokobori<sup>1</sup>, Akihiko Yamagishi<sup>1</sup>

1東京薬科大学大学院生命科学研究科

<sup>1</sup>Tokyo Univ. of Pharmacy and Life Science

Terrestrial microbes may be transported via various ways into the upper atmosphere. Due to the environmental similarity of the upper atmosphere to outer space, knowledge of microbiology in the upper atmosphere would be valuable for assessing the chances and limits of microbial transfer from the earth to extraterrestrial bodies (i.e., Panspermia of terrestrial microbes). We collected air dust samples in the upper troposphere and the stratosphere over Japan by using aircrafts or balloons. Microbial isolates from the samples were endospore-forming species (Bacillus, Paenibacillus, Streptomyces) and non-spore-forming Deinococci [1, 2]. Besides the evidence of microbial presence in the upper atmosphere, we show the possible presence of terrestrial microbes in space by height-dependent distribution of microbes [3]. The common characteristics of these upper atmospheric isolates were high resistance to radiation and desiccation, which likely enabled their survival in the environment of elevated radiation and desiccation. Specifically, the upper-atmospheric Deinococcus isolates exhibited extreme resistance to radiation (several times higher than bacterial endospores), the principle threat for microbial survival during interplanetary transfer. Therefore, Panspermia of viable Deinococci would be more likely than other terrestrial microbes [4]. Based on detailed characterization of the Deinococcus isolates, we proposed two new species Deinococcus aerius sp. nov. and Deinococcus aetherius sp. nov. [5, 6], which are now candidates for microbial exposure experiments in space.

 Y. Yang, S. Itahashi, S. Yokobori and A. Yamagishi, 2008. UV-resistant bacteria isolated from upper troposphere and lower stratosphere. Biological Sciences in Space 22 (1): 18-25.
Y. Yang et al., 2009. Investigation of cultivable microorganisms in the stratosphere collected by using a balloon in 2005. JAXA Research and Development Report, JAXA-RR-08-001: 35-42.
Y. Yang, S. Yokobori and A. Yamagishi, 2009. Assessing Panspermia Hypothesis by microorganisms collected from the high altitude atmosphere. Biological Sciences in Space 23 (3): 1 51-163.

4. Y. Yang, S. Yokobori and A. Yamagishi, 2009. Bacterial survival in response to desiccation and high humidity at above zero and subzero temperatures. Advances in Space Research, 43 (8): 1285 ?1290.

5. Y. Yang, T. Itoh, S. Yokobori, S. Itahashi, H. Shimada, K. Satoh, H. Ohba, I. Narumi and A. Yamagishi. 2009. Deinococcus aerius sp. nov., isolated from the high atmosphere. International Journal of Systematic and Evolutionary Microbiology 59 (8): 1862-1866.

6. Y. Yang, T. Itoh, S. Yokobori, H. Shimada, S. Itahashi, K. Satoh, H. Ohba, I. Narumi and A. Yamagishi, 2010. Deinococcus aetherius sp. nov., isolated from the stratosphere. International Journal of Systematic and Evolutionary Microbiology (in press).