

BBG006-02

会場:展示ホール7別室2

時間: 5月26日15:45-16:00

炭素・窒素安定同位体分析法を用いたサンゴ礁生態系における低次栄養段階の食物網の解明

Elucidating food webs of lower trophic levels in a coral reef ecosystem by carbon and nitrogen stable isotope analyses

中富 伸幸^{1*}, 中嶋 亮太¹, 吉田 輝明², 園田 和彦¹, 山本 修一¹, B.H.R. Othman², 戸田 龍樹¹

Nobuyuki Nakatomi^{1*}, Ryota Nakajima¹, Teruaki Yoshida², Kazuhiko Sonoda¹,
Shuichi Yamamoto¹, B.H.R. Othman², Tatsuki Toda¹

¹創価大学工学部, ²マレーシア国立大学 海洋生態研究センター

¹Faculty of Engineering, Soka University, ²Marine Ecosystem Research Center, UKM

The food webs of lower trophic levels in coral reef ecosystems have long been a black box in which secondary and fish productions are sustained being high, even though primary production is considerably low due to their oligotrophic environments. As possible alternative carbon sources in the ecosystems, several investigations suggested the importance of coral mucus and benthic microphyte. Yet, in order to quantitatively and qualitatively understand the flows of carbon and materials, it is necessary to make sure how those possible carbon sources are transferred from lower to higher trophic levels. To clarify what happens in such a black box, therefore, it is essential to look at the carbon sources and trophic structures of zooplankton communities that serve as energy and material transporters at lower trophic levels. We investigated carbon and nitrogen stable isotope of sorted zooplankton groups and possible alternative carbon sources. Zooplankton, coral mucus, suspended sediment and seawater samples were collected from the coral reef at Bidong Island, Malaysia in August 2009. Zooplanktons were taxonomically sorted whenever possible and copepods were specifically classified into genus-levels. We measured their abundance and biomass, as well as chlorophyll a concentrations of suspended sediment and seawater. The delta the ¹³C value of phytoplankton was estimated from the average values of herbivorous zooplankton groups. Their values ranged from -23 to -21 per mill and were fit into the range of the reported values in previous literature. The delta ¹³C and delta ¹⁵N values of 36 zooplankton groups ranged from -22.5 to -13.5 per mill and 2.4 to 7.3 per mill, respectively. They were statistically categorized into two clusters: one indicating the delta ¹³C values similar to the estimated values of phytoplankton (ca.-22.0 per mill), and another with the values close to coral mucus (-14.8 per mill). Also, most zooplankton groups including 15 genera of copepod had lower delta ¹³C (ca.-21.5 per mill), but several abundant groups with higher delta ¹⁵N such as mysids, fish larvae and siphonophora showed significantly higher delta ¹³C values (ca.-18.7 per mill) than those of the estimated food chain based on phytoplankton. These results suggest that although most groups can be categorize into the food chain originated from the carbon sources with lower delta ¹³C (most likely phytoplankton), the contribution of coral mucus that indicate higher delta ¹³C becomes more substantial as trophic levels increase in the coral reef ecosystem.

キーワード:安定同位体,サンゴ礁,動物プランクトン,食物網,低次栄養段階

Keywords: Stable isotope, coral reef, zooplankton, food web, lower trophic level