

BPT011-04

会場: 301B

時間: 5月24日16:15-16:30

## 白亜紀化石ココリスの結晶方位の多様性と共通性

### Crystallographic variation and uniformity in Cretaceous fossil heterococcoliths

猿渡 和子<sup>1\*</sup>, 田中裕一郎<sup>2</sup>, 長澤寛道<sup>3</sup>, 小暮敏博<sup>1</sup>

Kazuko Saruwatari<sup>1\*</sup>, Tanaka Yuichiro<sup>2</sup>, Nagasawa Hiromichi<sup>3</sup>, Kogure Toshihiro<sup>1</sup>

<sup>1</sup>東京大学大学院理学系研究科, <sup>2</sup>産業技術総合研究所, <sup>3</sup>東京大学大学院農学生命科学研究科

<sup>1</sup>U. Tokyo, Dept. EPS, <sup>2</sup>AIST, <sup>3</sup>U. Tokyo, Agriculture and Life science

Coccoliths are calcified scales formed around unicellular algae, coccolithophores, and they consist of radial arrays of calcite crystal units with complex morphologies. Since their first appearance at the Late Triassic (230 Ma), it is considered for the phyletic evolution that the morphological properties have variously changed but the crystallographic property has been conserved by the V/R model in which coccoliths are generally consist of two calcite units with sub-vertical (V-units) and sub-radial (R-units) c-axis (Young et al., 1992). However, the recent SEM-EBSD analyses revealed that a Cretaceous fossil coccolith, *Watznaueria barnesae*, has a different type of R-units compared with the living coccoliths, *Emiliana huxleyi* and *Gephyrocapsa oceanica*, suggesting the crystallographic variations of fossil coccoliths (Saruwatari et al., 2008). In order to elucidate the crystallographic variation of fossil coccoliths, we determined detailed crystallographic properties of several major genera of coccoliths in Cretaceous chalk from Dover, England, using electron backscattered diffraction (EBSD) analyses and SEM stereo-photogrammetry.

The results showed that both V- and R-units are divided into the several types with respect to the inclination and orientations of c- and a-axes, indicating the crystallographic variations of Cretaceous fossil coccoliths. As a general property, all units are likely aligned with the calcite  $\langle 4\ 8\ -1 \rangle$  direction parallel to coccolith rings. Concerning the morphology, most of the heterococcoliths developed calcite rhombohedral  $\{10\text{-}14\}$  surfaces but the imbricating murolith V-units did the  $\{2\text{-}1\text{-}10\}$  planes. These results indicate that crystallographic properties of both V- and R-units varied with their morphologies for the past 230 million years, but are regulated by the alignments of calcite  $\langle 4\ 8\ -1 \rangle$  directions along the coccolith ring.

キーワード: ヘテロココリス, 結晶方位, 方解石, 電子後方散乱回折, 白亜紀

Keywords: heterococcolith, crystallographic orientation, calcite, EBSD, Cretaceous