

BPT014-03

会場: 301B

時間: 5月27日09:30-09:45

## サンゴ礁域の大型有孔虫を用いた酸性化実験：石灰化率への影響検証

### Testing of the effect of ocean acidification on large benthic foraminifers

黒柳 あずみ<sup>1\*</sup>, 川幡 穂高<sup>1</sup>, 鈴木 淳<sup>2</sup>, 藤田 和彦<sup>3</sup>, 入江 貴博<sup>4</sup>

Azumi Kuroyanagi<sup>1\*</sup>, hodaka kawahata<sup>1</sup>, Atsushi Suzuki<sup>2</sup>, Kazuhiko Fujita<sup>3</sup>, Takahiro Irie<sup>4</sup>

<sup>1</sup>東京大学海洋研究所, <sup>2</sup>産業技術総合研究所地質情報研究部門, <sup>3</sup>琉球大学理学部物質地球科学科,  
<sup>4</sup>琉球大学熱帯生物圏研究センター

<sup>1</sup>Ocean Research Institute, Univ. of Tokyo, <sup>2</sup>Geological Survey of Japan, AIST,

<sup>3</sup>Faculty of Science, Univ. of the Ryukyus, <sup>4</sup>University of the Ryukyus

Modern surface seawater is saturated with respect to calcium carbonate, including calcite, high-Mg calcite (magnesian calcite), and aragonite. However, an increase in the atmospheric CO<sub>2</sub> partial pressure (pCO<sub>2</sub>) causes, via air-sea CO<sub>2</sub> exchange, the total dissolved CO<sub>2</sub> concentration in surface seawater to increase, and seawater pH to decrease. Ocean acidification has become recognized recently to be a major threat to calcifying organisms. Previous studies have reported that calcification rates of calcareous marine organisms (e.g., corals, foraminifers, coccolithophores, pteropods, mussels, and oysters) change in response to lowering pH levels even in waters oversaturated with respect to calcium carbonate. However, the impact of ocean acidification on large benthic foraminifers, which are major contributors to organic and inorganic carbon production in coral reefs, is still unclear. In this study, we cultured asexually-produced individuals of *Marginopora kudakajimensis* under four different pH conditions to examine the effects of lowering pH on their growth rates. Experimental results indicate that growth rate, measured by shell diameter, shell weight, and the number of chambers added, generally decreased with lowering pH after 10 weeks of culture. Shell weight was most closely dependent upon pH, suggesting that fossil shell weight can be more useful for reconstruction of paleo-pH changes. The relationship between the shell weight and shell size also showed significant differences among the four pH conditions. Between pH 8.3 and 7.7 (NBS scale), the tendency of the growth rate of *M. kudakajimensis* to decrease with pH was consistent with that of most other calcifying organisms. However, the calcification/growth rates at pH 7.9 and ~8.2 (present seawater value, control) were not significantly different, and other organisms also display a nonlinear response to elevated pCO<sub>2</sub> at around this pH range. These results suggest that 1) they already may have experienced a reduction in growth in natural environments since preindustrial times and 2) although the seawater CO<sub>2</sub> system of reef water shows great variation, the calcification rate of these large foraminifers should remain at the present level at pH 7.9-8.2. However, at around pH 7.7, their calcification rate would decline steeply, which would probably preclude their survival.

キーワード: 大型有孔虫, 海洋酸性化, 飼育実験, 石灰化率

Keywords: larger benthic foraminifer, ocean acidification, culture experiment, calcification rate