

BPT014-P04

会場:コンベンションホール

時間: 5月27日17:15-18:45

石垣島の津波石ハマサンゴを用いた古環境復元とジオハザードへの応用

Reconstruction of paleoenvironment and paleo-geohazard recorded in Porites coral boulders cast ashore in Ishigaki Island

荒岡 大輔^{1*}, 井上 麻夕里², 鈴木 淳³, 横山 祐典², ラリー エドワーズ⁴, ハイ チェン⁴, 後藤 和久⁵, 河名 俊男⁶, 松崎 浩之⁷, 菅 浩伸⁸, 鹿園 直建⁹, 川幡 穂高¹

Daisuke Araoka^{1*}, Mayuri Inoue², Atsushi Suzuki³, Yusuke Yokoyama², R. Lawrence Edwards⁴, Hai Cheng⁴, Kazuhisa Goto⁵, Toshio Kawana⁶, Hiroyuki Matsuzaki⁷, Hironobu Kan⁸, Naotatsu Shikazono⁹, hodaka kawahata¹

¹東大・院・新領域, 東大・海洋研, ²東大・海洋研, ³産総研・地質情報, ⁴ミネソタ大学,
⁵東北大・工・災害制御研究センター, ⁶琉大・教育, ⁷東大・院・工学, ⁸岡山大・教育, ⁹慶應・理工

¹GSFS and ORI, The Univ. of Tokyo, ²ORI, The Univ. of Tokyo, ³GSI, AIST, ⁴Univ. of Minnesota,
⁵DCRC, Tohoku Univ., ⁶Univ. of the Ryukyus, ⁷MALT, The Univ. of Tokyo, ⁸Okayama Univ., ⁹Keio Univ.

A large number of massive coral boulders are scattered on the shore and in the reef of eastern Ishigaki Island, Japan. Many of these boulders were likely cast ashore by the 1771 Meiwa earthquake tsunami, which was one of the largest tsunami disasters in Japanese history, resulting in the deaths of 12,000 people in the region. According to the previous study, ¹⁴C ages of well-preserved surface parts of these coral boulders, particularly focused on Porites spp. corals, were analyzed with the aim of confirming that they had been transported by the Meiwa tsunami. However, the calibrated ¹⁴C age distribution showed a relatively wide range, and the events transporting these boulders onshore could not be entirely clear, although it was revealed that massive Porites coral boulders had not been caused by a single event.

Precise and accurate dating of coral sample is now possible by ²³⁰Th dating. To confirm when coral boulders had been cast ashore and dead, we applied high-precision ²³⁰Th dating to well-preserved surface parts of massive Porites coral boulders.

Coral skeletons are composed of calcium carbonate, and the stable isotope ratios and trace elements in the skeleton can provide insight into environmental information. Therefore, modern and fossil Porites corals have been used for paleoenvironmental reconstructions to reveal past oceanographic and climatic changes. By using ²³⁰Th dating for Porites coral boulders and splicing paleoenvironmental data reconstructed from coral skeletons of several these boulders dying at various dates, it has a potential to reconstruct paleoenvironment at high-resolution for several centuries, continuously.

Another aim of this study is to determine the ages of past tsunamis and/or severe storm events by high-precision ²³⁰Th dating of youngest part of Porites coral boulders. In addition, identifying coral boulders cast ashore by the 1771 Meiwa tsunami has scientific merit for constraining their tsunami models to apply these boulders. It is important to determine the seismic recurrence period for long-term assessment of future earthquake and tsunami risk in this region, and, if the boulders were cast onshore by large typhoons, the frequency of such events should also be known for future disaster mitigation. Our study demonstrated that Porites boulders are useful for studies of not only paleoenvironment but also paleo-geohazard such as tsunamis and extreme storms.

キーワード:ハマサンゴ,津波石, 230Th年代測定法, 14C年代測定法,ジオハザード,古環境

Keywords: Porites sp. coral, Tsunami boulder, 230Th dating, 14C dating, paleo-geohazard, paleoenvironment