## **Japan Geoscience Union Meeting 2010**

(May 23-28 2010 at Makuhari, Chiba, Japan)

©2009. Japan Geoscience Union. All Rights Reserved.



MIS001-04 会場: 202

時間: 5月27日11:30-11:45

A statistical study on the correlation between lower ionospheric perturbations as seen by subionospheric VLF/LF propagat

A statistical study on the correlation between lower ionospheric perturbations as seen by subionospheric VLF/LF propagat

早川 正士<sup>1\*</sup>, Y. Kasahara<sup>2</sup>, T. Nakamura<sup>2</sup>, F. Muto<sup>2</sup>, T. Horie<sup>2</sup>, S. Maekawa<sup>2</sup>, Y. Hobara<sup>2</sup>, A.A. Rozhnoi<sup>3</sup>, M. Solovieva<sup>3</sup>, O.A. Molchanov<sup>3</sup>

Masashi Hayakawa<sup>1\*</sup>, Y. Kasahara<sup>2</sup>, T. Nakamura<sup>2</sup>, F. Muto<sup>2</sup>, T. Horie<sup>2</sup>, S. Maekawa<sup>2</sup>, Y. Hobara<sup>2</sup>, A.A. Rozhnoi<sup>3</sup>, M. Solovieva<sup>3</sup>, O.A. Molchanov<sup>3</sup>

¹電気通信大学 AWCC, ²電気通信大学, ³Institute of Physics of the Earth

<sup>1</sup>UEC AWCC, <sup>2</sup>UEC, <sup>3</sup>Institute of Physics of the Earth

The subionospheric VLF/LF propagation is extensively used to study the lower ionospheric perturbation in possible association with earthquakes. An extensive period of data over seven years and a combination of different propagation paths are used to examine the statistical correlation between the VLF/LF propagation anomaly (average nighttime amplitude, dispersion and nighttime fluctuation) and earthquakes with magnitude greater than 6.0. It is then found that the propagation anomaly exceeding the 2 sigma (standard deviation) criterion indicating the presence of ionospheric perturbation is significantly correlated with earthquakes with shallow depth (<40km). Finally some comments on the mechanism of seismo-ionospheric perturbations are discussed.

Keywords: VLF/LF propagation, earthquakes, statistical study