

## モンゴルのジュラ系-白亜系湖成層を対象とした陸域古環境変遷の高解像度復元

### Reconstruction of terrestrial paleoenvironmental change from the Jurassic-Cretaceous lacustrine deposits in SE Mongolia

長谷川 精<sup>1\*</sup>, 安藤 寿男<sup>2</sup>, 長谷川 卓<sup>3</sup>, 太田亨<sup>4</sup>, 山本正伸<sup>5</sup>, 長谷部 徳子<sup>6</sup>, ガン・リー<sup>7</sup>, ニーデン・イチノロフ<sup>8</sup>

Hitoshi Hasegawa<sup>1\*</sup>, Hisao Ando<sup>2</sup>, Takashi Hasegawa<sup>3</sup>, Tohru Ohta<sup>4</sup>, Masanobu Yamamoto<sup>5</sup>, Noriko Hasebe<sup>6</sup>, Gang Li<sup>7</sup>, Niiden Ichinnorov<sup>8</sup>

<sup>1</sup>北海道大学創成研究機構, <sup>2</sup>茨城大学理学部, <sup>3</sup>金沢大学理学部, <sup>4</sup>早稲田大学教育学部,

<sup>5</sup>北海道大学地球環境科学研究所, <sup>6</sup>金沢大学環日本海域環境研究センター, <sup>7</sup>南京地質古生物研究所,

<sup>8</sup>モンゴル科学アカデミー, 古生物センター

<sup>1</sup>CRIS, Hokkaido University, <sup>2</sup>Dept. of Earth Sciences, Ibaraki Univ., <sup>3</sup>Dept. of Earth Sciences, Kanazawa Univ.,

<sup>4</sup>Faculty of Edu., Int. Arts, Sci., Waseda, <sup>5</sup>Faculty of Env. Ear. Sci., Hokkaido Univ,

<sup>6</sup>Division of Ear., Env. Sci., Kanazawa U., <sup>7</sup>Nanjing Inst. of Geol. and Palaeont., <sup>8</sup>Paleontol. Center, MAS, Mongolia

The Jurassic-Cretaceous period is characterized by an extremely warm "greenhouse" climate, elevated atmospheric CO<sub>2</sub> levels, and repeated occurrences of Ocean Anoxic Events (OAEs); however, detailed processes and causal mechanisms of these marked events, particularly the response of terrestrial climate system, have been poorly understood. Possible causal mechanisms of OAEs in the greenhouse conditions include enhanced surface productivity and the excess of organic burial in the oceans. Terrestrial weathering may have increased terrigenous input into the oceans ("Weathering Hypothesis"; e.g., Weisert et al., 1998; Hasegawa, T., 2003). To evaluate interaction between the land and the ocean during OAE interval, we investigated terrestrial paleoenvironments at intra-continental sites in eastern Eurasia.

We examined the lower Cretaceous and the middle-late Jurassic lacustrine deposits (Shinekhudag and Khamarkhoovor formations, respectively) in southeast Mongolia. The Shinekhudag Formation, well exposed in the Shine Khudag locality, is composed mainly of paper shale, oil shale and silty mudstone, with high content of organic carbon. Strata are continuously exposed about 500 m in thickness. The age of the Shinekhudag Formation is estimated as Aptian or Barremian-Aptian based on the floral and molluscan evidence (Martinson and Shuvalov, 1973; Krassilov, 1982; Jerzykiewicz and Russell, 1991), and K-Ar age data of basaltic rocks in the uppermost part of the underlying Tsagantsav Formation (ca. 126 Ma: Graham et al., 2001). Khamarkhoovor Formation, well exposed in the Eedemt locality, is also composed of paper shale, oil shale and silty mudstone, with higher content of organic carbon. The age of the strata in Eedemt locality is newly defined as middle-late Jurassic, based on the conchostracans biostratigraphy (Li et al., in submitted). These shale and silty mudstone successions are rhythmically alternated (decimeter-, meter-, tens of meter-scale) both in Shine Khudag and Eedemt localities, which can be controlled by orbital cycles.

To establish detail chronostratigraphy of the middle Jurassic-lower Cretaceous lacustrine deposits and reconstruct terrestrial paleoenvironmental changes, we are conducting the following

approaches; 1) carbon-isotope stratigraphy; 2) cyclostratigraphy of the rhythmically alternated shale and silty mudstone successions; 3) conchostracans biostratigraphy; 4) fission-track age dating of intercalated tuff; 5) reconstruction of paleoweathering changes based on the major elemental ratios and weathering index (Ohta & Arai, 2007); and 6) paleotemperature reconstruction with organic proxies (e.g., TEX86). Establishment of the carbon-isotope stratigraphy and high resolution cyclostratigraphy enable us a high-resolution and precise correlation of the terrestrial paleoenvironmental changes with the specific global events (such as OAEs). Understanding of the global climate system, in particular, the terrestrial climate during the past "greenhouse" period gives clue to future prediction of anthropogenic global warming.

キーワード:湖成層,白亜紀,ジュラ紀,陸成,サイクル層序,海洋無酸素事変

Keywords: lacustrine deposits, Cretaceous, Jurassic, terrestrial, cyclostratigraphy, OAEs