

Effect of Carbon and Hydrogen on Melting and Magma properties at High Pressure

Eiji Ohtani^{1*}, Tatsuya Sakamaki¹, Konstantin Litasov¹

¹Tohoku University

CO₂ and H₂O are important components in the Earth's mantle. These species have strong effects on melting and melt properties such as density and viscosity under the mantle conditions. We studied the effect of CO₂ and H₂O on melting relations of the mantle materials up to 30 GPa. The experiments revealed that the solidus temperature in the peridotite-CO₂ system is comparable with that of the wet solidus of peridotite in the deep upper mantle, whereas it is lower than the wet solidus under the transition zone and lower mantle conditions. Thus, the fluid enriched in CO₂ is likely to be formed under the transition zone and lower mantle conditions, whereas hydrous fluid can be formed more easily under the upper mantle conditions.

The density of the volatile rich magmas is controlled by the partial molar volume of the volatile components in magmas. We have determined the partial molar volume of H₂O and CO₂ in magmas at high pressure up to about 20 GPa by a sink-float experiment using a diamond density marker and the X-ray absorption method. Our measurements revealed that the CO₂ component dissolved in magmas is less compressible compared to the H₂O component at high pressure. These compression behaviors of CO₂ and H₂O components in magmas indicate that CO₂ reduces the magma density more effectively compared to H₂O, and density crossover at the base of the upper mantle between volatile-rich magmas and surrounding peridotite mantle is less likely if the volatile is CO₂ enriched in compositions, since CO₂ can reduce the magma density effectively even at high pressure of the base of the upper mantle. The CO₂ rich melts formed in the upper mantle, transition zone and lower mantle can be buoyant and can move upwards.

Keywords: carbon, hydrogen, melting, magma, density, fluid