

中国雲南省三江地域に分布する変成岩類の変成作用

Metamorphism of the Sanjiang region in Yunnan province, SW China.

米村 和紘^{1*}, 小山内 康人¹, 中野 伸彦¹, 大和田 正明², 馬場 壮太郎³, 加々島 慎一⁴,
野原 (今中) 里華子⁵

Kazuhiro Yonemura^{1*}, Yasuhito Osanai¹, Nobuhiko Nakano¹, Masaaki Owada², Sotaro Baba³,
Shin-ichi Kagashima⁴, Rikako Nohara-Imanaka⁵

¹九大・比文・地球変動, ²山口大・理・地球科学, ³琉大・教・自然環境, ⁴山形大・理・地球環境,
⁵新潟大・自然科学

¹Earth Sci., Kyushu Univ, ²Dept. Earth Sci., Yamaguchi Univ., ³Natural Environment, Univ. of Ryukyus,
⁴Earth&Environmental Sci., Yamagata Univ., ⁵Grad. Sch. Sci. & Tech., Niigata Univ.

Several micro-continents had been collided in the Sanjiang region in Yunnan province, SW China. We studied two areas geologically and petrologically, Cangshan area in the east and Nujiang area in the west. The Cangshan area is considered to represent a possible northern extension of the Ailao Shan - Red River shear zone, the collision boundary between South China craton and Indochina craton. The Nujiang area is also considered as the collision boundary between Indochina craton and Shan-Thai craton. Detailed studies of the metamorphic rocks from the both areas give us useful information to understand the tectono-metamorphic evolution during the Asian continental growth.

The metamorphic rocks from the Cangshan area demonstrate greenschist- to upper amphibolite-facies conditions. The mineral assemblages of metamorphic rocks changes systematically from east to west. Based on the change in mineral assemblages of the pelitic to psammitic rocks, the Cangshan area can be divided into the following four metamorphic zones; zone A Chl + Ms in the east, zone B Grt + Bt + Ms and Grt + St + Bt + Ms, zone C Grt + Sil + Bt, zone D Chl + Ms in the west. The estimated P-T condition of zone C is 800-850 C, 10-11 kbar from Ky-bg Grt-Sil-Bt gneiss. Reaction textures and chemical zoning of garnet suggest a clockwise P-T path, and the geothermal gradient suggests a medium pressure type.

The metamorphic rocks from the Nujiang area suggest greenschist- to amphibolite-facies conditions. The mineral assemblages of metamorphic rocks changes systematically from west to east. Metamorphic grade of mafic rocks also grade into the east. Based on the changes in mineral assemblages of the metapelites, the Nujiang area can be subdivided into the following five metamorphic zones; zone I Chl + Ms in the west, zone II Grt + And + Bt + Ms, zone III Grt + Sil + Bt, zone IV Grt + Crd + Bt, zone V Ms in the east. The estimated P-T conditions are 700-830 C, 4-6 kbar (Grt-Sil-Crd-Bt gneiss) and 650-780 C (Grt-Crd-Bt gneiss) in the zone IV. Several metamorphic reaction textures suggest heating process at low pressure condition, and the metamorphic field gradient is confirmed as a low pressure type.

In the Cangshan area, the chondrite normalized REE patterns show three different patterns; LREE enriched pattern, flat pattern like E-MORB and depleted REE pattern with Eu positive anomaly. The epsilon Sr (eSr) values, normalized at 250Ma, are from -13.57 to 127.78 and epsilon Nd (eNd) values are from -8.04 to 5.84, respectively. They are divided into high-NdI group and low-NdI

group. High-NdI group is similar to mafic rocks from the Red River area. Low-NdI group suggests a cumulate origin because it shows depleted REE pattern. In the Nujiang area, the chondrite normalized REE patterns also indicate two patterns; LREE enriched pattern and flat pattern like E-MORB. The ϵ_{Sr} values, normalized 250Ma, are from -1.90 to 139.17 and ϵ_{Nd} values are from -1.68 to 4.33. In the Red River area, where's tectonics related to the Cangshan area, the chondrite normalized REE patterns indicate two different patterns; LREE enriched pattern and flat pattern like E-MORB. The ϵ_{Sr} values, normalized 250Ma, are from 2.60 to 49.60 and ϵ_{Nd} values are from -0.41 to 3.18.

Estimated medium pressure geothermal gradients and geochemical characteristics of the metamorphic rocks from the Cangshan area are similar to those of the Red River area. These similarities suggest both areas related to the same collision tectonics. On the other hand, metamorphic rocks from the Nujiang area were formed by simple heating process under the upper crustal condition. The differences of metamorphism indicate that the Sanjiang region formed by multi-stage collision tectonic events during the growth of Asian continent.