

SCG084-P12

Room: Convention Hall

Time: May 25 17:15-18:45

## Application of groundwater dating in Horonobe area, Hokkaido, Japan

Kouki Kashiwaya<sup>1\*</sup>, Takuma Hasegawa<sup>1</sup>, Kotaro Nakata<sup>1</sup>, Takahiro Oyama<sup>1</sup>, Yuichi Tomioka<sup>1</sup>

<sup>1</sup>CRIEPI

Knowledge on groundwater flow property around disposal site is essential for safety assessment of radioactive waste disposal. Central Research Institute of Electric Power Industry (CRIEPI) has investigated groundwater dating methods such as <sup>4</sup>He accumulation method and <sup>36</sup>Cl method that enable estimation of very long residence time.

The <sup>4</sup>He accumulation method can estimate residence time based on increment of <sup>4</sup>He concentration in groundwater and generation rate of <sup>4</sup>He in rock. <sup>36</sup>Cl is a radioisotope whose half-life period is 301 thousand years, and the <sup>36</sup>Cl method is based on concentration change resulted from <sup>36</sup>Cl decay in groundwater. Applicability of these methods was confirmed in previous works carried out in the Great Artesian Basin, Australia.

Research using these methods was conducted in Horonobe, Hokkaido, Japan, to consider the applicability of the methods and to adapt them to geological environment in Japan. Neogene sedimentary formations, Wakkanai and Koitoi formations are thickly distributed around the area, and the groundwater dating was carried out with cores and groundwater collected from the formations.

<sup>4</sup>He concentration in the samples was in the range of  $10^{-6}$ - $10^{-5}$  cc<sub>STP</sub>/g<sub>w</sub>. Equilibrium concentration of <sup>4</sup>He in atmosphere is  $4.8 \times 10^{-8}$  cc<sub>STP</sub>/g<sub>w</sub> and generation rate of <sup>4</sup>He in situ is  $1-2 \times 10^{-12}$  cc<sub>STP</sub>/g<sub>w</sub>y and thus residence time of groundwater was calculated as several million years. However, <sup>3</sup>He/<sup>4</sup>He ratio of He in groundwater was in the range of  $10^{-7}$ - $10^{-6}$  and slightly high relative to the <sup>4</sup>He concentration. This may imply the inflow of He from external origin.

<sup>36</sup>Cl/Cl of the samples was in the range of  $10^{-15}$ - $10^{-14}$ . The origin of deep groundwater in Horonobe area is considered to be seawater trapped when the formations were formed. Averaged <sup>36</sup>Cl/Cl of present seawater obtained in previous works was  $1.0 \times 10^{-15}$ , and the <sup>36</sup>Cl/Cl values of groundwater was higher. This means that <sup>36</sup>Cl/Cl in groundwater increased by radioactivation in the formations. Considering the half-life period of <sup>36</sup>Cl, residence time was estimated as larger than 1.5 million years. The residence time obtained from the two methods showed that groundwater flow in this area is not active.

### Acknowledgements

This research entitled 'Research and development on groundwater dating technique' was carried out under contracts awarded by the Japanese Ministry of Economy, Trade and Industry (METI) to CRIEPI. In-situ investigations in Horonobe study site have been conducted as joint research with Japan Atomic Energy Agency.

Keywords: groundwater, residence time, sedimentary rock, groundwater dating