

SCG085-05

会場: 201B

時間: 5月23日16:22-16:35

コーダ波干渉法を用いた離散的深部低周波微動間の距離決定の試み

Attempt to obtain separations of discrete deep low frequency tremor with coda wave interferometry

宮澤 理稔<sup>1\*</sup>, 太田 和晃<sup>2</sup>, 井出 哲<sup>2</sup>

Masatoshi Miyazawa<sup>1\*</sup>, Kazuaki Ohta<sup>2</sup>, Satoshi Ide<sup>2</sup>

<sup>1</sup>東大・地震研, <sup>2</sup>東大・理

<sup>1</sup>ERI, Univ. Tokyo, <sup>2</sup>Grad. School Sci., Univ. Tokyo

We attempt to obtain separations between discrete episodes of deep low frequency (DLF) tremor from the coda waves. Since the DLF tremor has emergent arrivals, it is still difficult to pick P and S wave arrivals to obtain the source location and constrain the separations. Assuming each DLF tremor has a similar mechanism and occurs close to one another, we use the coda parts of the observed wave trains and apply coda wave interferometry (Snieder and Vrijlandt, 2005). The result shows that the DLF tremor sequence occur within an interval of about 100 m when the each discrete episode has duration of about 10 s.

キーワード: 深部低周波微動, コーダ波干渉法

Keywords: deep low frequency tremor, coda wave interferometry