

小笠原海台および母島海山に分布する炭酸塩岩の堆積相・年代とその地質学的意義

Cretaceous shallow-water carbonates from the Hahajima Seamount and the Ogasawara Plateau

二宮 淳¹, 高柳 栄子¹, 石川 剛志², 永石 一弥³, 西村昭⁴, 中澤努⁴, 柴崎 洋志⁵, 井龍 康文^{1*}

Jun Ninomiya¹, Hideko Takayanagi¹, Tsuyoshi Ishikawa², Kazuya Nagaishi³, Akira Nishimura⁴, Tsutomu Nakazawa⁴, Hiroshi Shibasaki⁵, Yasufumi Iryu^{1*}

¹名古屋大学大学院環境学研究科, ²海洋研究開発機構高知コア研, ³マリン・ワーク・ジャパン, ⁴産業技術総合研究所, ⁵石油天然ガス・金属鉱物資源機構

¹Nagoya Univ., ²JAMSTEC, ³Marine Works Japan, ⁴AIST, ⁵JOGMEC

The Hahajima Seamount is situated at the junction between the Izu-Bonin and Mariana forearc slopes and is located at a collision area between the forearc slope of the Philippine Sea Plate and the Ogasawara Plateau on the Pacific Plate. Shallow-water carbonates were recovered from the eastern slope of this seamount. Five seamounts (Katayama, Hanzawa, Yabe, Higashi, and Minami Seamounts), which are capped with thick sequences of shallow-water carbonates, are located on the Ogasawara Plateau.

The shallow-water carbonates on the Hahajima Seamount and the Ogasawara Plateau are similar in lithology and age. They are dominated by molluscan floatstone/rudstone with common intraclasts. Gravel-sized bioclasts include molluscs (e.g., *Nerinea* sp. and rudists) and, locally, calcareous sponges and minor amounts of coral. Sand-sized components include bioclasts of benthic foraminifers and dasycladacean algae and non-skeletal grains of intraclasts and peloids, generally with few ooids. Most of the bioclasts are bored and micritized and frequently have thick micrite envelopes. Their Sr isotope ages fall into two ranges, 112.3-113.6 Ma (Aptian) and 76.4-88.7 Ma (Coniacian to Campanian). These similarities in lithology and Sr isotope ages, as well as the limited occurrence of carbonates on the Hahajima Seamount, suggest that they were not deposited in situ but originated from the Ogasawara Plateau. The extensive bioerosion may indicate that a relatively high nutrient level existed in the tropical shallow marine environment during the Cretaceous.

The current reconstruction of plate motion based on magnetic anomalies of the ocean floors suggests that the oceanic crust on which the Ogasawara Plateau lies formed in the late Jurassic (about 153 Ma) at 20 degrees south latitude. Then, the plateau moved west-northwestward until 100 Ma, when carbonate platforms formed at about 15 degrees south latitude (Aptian).

Subsequently, the plate moved north-northeastward from about 100-90 Ma (Albian-Turonian) and northward/ north-northwestward from about 90-80 Ma (Turonian-Campanian), during which time carbonate platforms extended from about 4-13 degrees north latitude on the plateau (Coniacian to Campanian).

Keywords: Hahajima Seamount, Ogasawara Plateau, shallow-water carbonate, Sr isotope age, Cretaceous