

SCG089-08

会場: 101

時間: 5月27日14:00-14:15

小笠原群島における島弧初期マグマの地球化学的特徴および起源について

Geochemical characteristics and the origin of the incipient arc magmas in the Bonin Islands

金山 恭子^{1*}, 海野 進², 石塚 治³

Kyoko Kanayama^{1*}, Susumu Umino², Osamu Ishizuka³

¹金沢大学大学院自然科学研究科, ²金沢大学理工学域, ³産業技術総合研究所地質情報研究部門

¹Graduate School of Kanazawa University, ²Kanazawa University, ³Geological Survey of Japan/AIST

Bonin Islands are known for the occurrence of boninite series and high-Mg arc tholeiite and calc-alkaline rock series generated at the incipient stage of formation of subduction zone. We present new analysis of major and trace elements and petrogenetic processes of volcanic rocks of the Bonin Islands. Boninite series rocks in Chichijima and Mukojima Island Group represent the primitive arc magmatism in middle Eocene time, which gave way to arc tholeiite and calc-alkaline rocks in Hahajima Island Groups. Boninite series of the Maruberiwan and Asahiyama Formations indicates a differentiation trend that sharply increases in FeO^*/MgO (0.47-32) with increasing SiO_2 contents (53.8-78.2 wt %). FeO^*/MgO ratios (0.74-4.19) of arc tholeiite and calc-alkaline rocks of the Hahajima Island Group are slightly lower than those of the Izu-Bonin Quaternary volcanic front lavas. Boninite series samples of the Mikazukiyama Formation show a similar trend to the Maruberiwan-Asahiyama boninite series samples. However, the former has lower SiO_2 contents (55.7-58.2 wt %), and higher FeO^*/MgO ratios (1.05-1.98) than the latter. Most of the Maruberiwan and Mikazukiyama boninites belong to low-Ca type, while only a small number of Maruberiwan boninite samples belong to high-Ca type.

Maruberiwan boninites ($\text{MgO} > 8$ wt%) are the most depleted in REEs ($\text{Yb} = 0.36\text{-}1.01$ ppm), which is only about 1/10 of N-MORBs, and the most enriched in LILEs ($\text{Rb} = 6\text{-}23$, $\text{Ba} = 6\text{-}58$ ppm) in the Bonin Islands. Low-Ca boninites indicate a distinct positive anomaly of Zr and negative anomalies of Sm and Ti ($(\text{Sm}/\text{Zr})_n = 0.45\text{-}0.78$, $(\text{Ti}/\text{Zr})_n = 0.25\text{-}0.64$), while high-Ca boninites indicate moderate anomalies of these elements ($(\text{Sm}/\text{Zr})_n = 0.79\text{-}1.29$, $(\text{Ti}/\text{Zr})_n = 0.57\text{-}1.02$). Basalts of the Hahajima Island Group are the most enriched in REEs ($\text{Yb} = 1.2\text{-}2.4$ ppm) and HFSEs ($\text{Nb} = 0.47\text{-}1.75$ ppm) in the Bonin Islands. Compared to the Hahajima basalts, the present Izu-Bonin front lavas are more depleted in LREEs and HFSE, and enriched in LILEs.

Estimated conditions of magma generation show that the primary magmas in the Hahajima Island Group can coexist with lherzolite residue at 1-1.3 GPa under dry or wet (water content < 0.5 %) condition.

Results of model calculations strongly suggest that the source mantle changed from highly depleted to fertile compositions together with slab-derived components that changed from silicate melt to aqueous fluid as the generated magma changed from boninite in shallow (~ 0.8 GPa) mantle 48 Ma to basalt in deeper (1-1.3 GPa) mantle 44 Ma. The differences in trace element compositions between low-Ca and high-Ca boninites originated from the slab component, but not the source depletion.

キーワード:小笠原群島,ボニナイト,玄武岩,微量元素,主要元素

Keywords: Bonin Islands, boninite, basalt, trace elements, major elements