

SEM032-03

会場:展示ホール7別室2

時間: 5月25日09:30-09:45

日本の広域テフラ中の磁性鉱物の岩石磁気学的同定

Rock magnetic identification of magnetic minerals in widespread tephra layers in Japan

鳥居 雅之^{1*}, 中原 佑正¹, 藤井 純子², 中島 正志², 山本 裕二³, 小玉 一人³

Masayuki Torii^{1*}, Yuhsei Nakahara¹, Junko Fujii², Tadashi Nakajima², Yuhji Yamamoto³, Kazuto Kodama³

¹岡山理科大学生物地球システム学科, ²福井大学教育地域科学部, ³高知大学海洋コア総合研究センター

¹Okayama University of Science, ²Fukui University, ³Kochi University

As a typical volcanic island, widespread tephra layers are ubiquitously found in various types of Japanese Quaternary sediments. Those tephra layers are having primary importance in correlation and age estimation among Quaternary sediments, where the identification of each tephra has been mainly performed based on refractive index of glass shards and heavy mineral composition. A few rock magnetic attempts in identifying tephra layer have been done, although the results were not always straightforward. Recently, highly comprehensive rock magnetic analysis was applied to an Alaskan tephra by Lacroix et al. (2004). They succeeded to reveal the coexistence of titanohematite and titanomagnetite with low-temperature magnetic methods and appealed an effectiveness of the rock magnetic method.

We applied rock magnetic methods to the several representative tephra layers. High-temperature and low-temperature magnetic measurements and IRM analysis were applied to the samples from following six tephra layers: K-Ah (c. 7 ka), AT (c. 29 ka), DKP (c. 55 ka), Aso4 (c. 90 ka), Ata (c. 110 ka), and SK (c. 115 ka), collected more than 100 localities (e.g., Nakajima and Fujii, 1995). The high-temperature measurement was done with a thermomagnetic balance at KCC to show the Curie temperature (T_c). MPMS was used to show temperature dependence of the low-temperature IRM at OUS with zero-field cooling and field cooling modes. MPMS was also used to give IRMs at 300 K from 1 mT to 5T at 100 points along a logarithmic axis. The IRM acquisition curve was then analyzed with Irmunmix V2.2 (Heslop et al., 2002).

All samples suggest the presence of more than two magnetic constituents. The Curie temperatures were observed at 300-400°C and 500-600°C. While high-temperature tail above 600°C is sometimes observed, only a few samples indicate T_c around 200°C. The Verwey transition at about 120 K was not always observed, which suggests rather uncommon bearing of stoichiometric magnetite in tephra. As tephra are generally quenched material, exsolution of magnetite through high-temperature oxidation could be restrained during eruption. Large decrease in IRM was sometimes observed at about 50 K, which suggests an occurrence of titanomagnetite. IRM analyses imply two or more coercivity components. It is noteworthy that extremely high coercive component (>1 T) is often estimated. As goethite is not indicated by thermomagnetic curves, titanohematite can be an alternative answer. These lines of evidences suggest the presence of titanomagnetites with minor titanohematite in the studied tephra.

キーワード:広域テフラ,磁性鉱物,岩石磁気学

Keywords: widespread tephra, magnetic minerals, rock magnetism