

Preliminary Results from the 2D Ocean Bottom Magnetotelluric modeling beneath the Marmara Sea, Turkey

Tulay Kaya^{1*}, Yasuo Ogawa¹, Takafumi Kasaya², Bulent Tank¹, M. Kemal Tuncer³, Naoto Oshiman⁴, Yoshimori Honkura¹, Masaki Matsushima¹

¹Tokyo Institute of Technology, ²JAMSTEC, ³Bogazici University, ⁴Kyoto University

Turkey is a natural laboratory for geoscientists as one of the seismically most active countries in the world. The relative motions of surrounding tectonic plates (Eurasian, African, and Arabian plates), and Anatolian block has caused the formation of the North and East Anatolian fault zones (EAFZ and NAFZ) which are two major intercontinental strike-slip faults of Turkey. Both NAFZ and EAFZ have produced several destructive earthquakes in the country over the years. NAFZ is a ~1600 km plate boundary between Eurasia and the Anatolian plate. Observed westward movement of the Anatolian plate with respect to stable Euroasia is considered to be main reason for the major events along this transform zone. Since 1939, the westward migration of the big earthquakes on NAF has become an important phenomenon and attracted the attention of scientific community. After the last two earthquakes which occurred at the eastern end of Marmara Sea in 1999 (Mw7.4 Izmit, Mw7.2 Duzce), the westward migration of earthquake sequence indicates that next earthquake can possibly be located inside the Marmara Sea. From the east to the west, the NAFZ zone is divided into several branches on land. However, our knowledge of the geometry of the Marmara branch of NAFZ, a crucial parameter for the magnitude predictions of next event, is insufficient. The crustal structure beneath the Marmara Sea is not well uncovered yet. In this study, our primary task is to reveal the crustal resistivity distribution and its relation with the fault network beneath the Marmara Sea. The previous on-shore and off-shore magnetotelluric studies show that there is a strong relation between the resistive areas and asperity zones. Most of the resistor-conductor boundaries coincide with fault zones. In this study, the resistivity structure in the Marmara Sea was investigated by the magnetotelluric method based on the ocean bottom electromagnetic (OBEM) data. Totally at 17 sites, continues electric and magnetic fields (three components) were recorded with a sampling rate of 8Hz during ~ 1 month along three profiles crossing NAF within the Marmara Sea. The time series analysis is achieved by the Chave and Thompson code (1989). We perform 2D inversion modeling using Ogawa and Uchida (1996) code modified by us to account for the bathymetry effects on the measurements.

Keywords: Marmara Sea, Magnetotelluric, Ocean Bottom Electromagnetic (OBEM), North Anatolian Fault (NAFZ), Electromagnetic, Conductivity