

SGC065-04

会場: 301A

時間: 5月23日09:45-10:00

Micro-XANESを用いた下部マントルの酸化還元状態の推定

Oxidation state of the lower-mantle revealed by micro-XANES measurements

小竹 翔子^{1*}, 鍵 裕之¹, 石橋 秀巳¹, Harte Ben²

Shoko Odake^{1*}, Hiroyuki Kagi¹, Hidemi Ishibashi¹, Harte Ben²

¹東大院・理・地殻化学, ²エジンバラ大

¹Geochem. Lab., Grad. School Sci. Univ., ²GeoScience, Edinburgh Univ.

The valence state of iron in mantle-derived minerals has been investigated as an indicator of redox condition of the earth's interior. However, it was reported that the valence state of iron in lower-mantle minerals is controlled by the crystallographic structure of the host minerals and is not sensitive to the redox conditions of the lower-mantle. Alternatively, we focus our attention on chromium. Chromium is mostly found as Cr³⁺ and Cr⁶⁺ in terrestrial minerals. However, minerals formed in extremely reduced conditions can contain Cr²⁺ in their crystal structure. Thus, the presence of Cr²⁺ in a mineral could be an indicator of the reduced environment. Diamonds which contain the lower-mantle minerals as inclusions are the deepest sourced materials. They are the only source which can provide direct information about the lower-mantle. In this study, we investigated the oxidation state of chromium in natural lower-mantle mineral, ferropericlasite, using XANES spectroscopy.

Chromium and Iron K-edge X-ray absorption near-edge structure (XANES) spectra were obtained on three natural ferropericlasite inclusions. These inclusions were picked up from diamonds originated from the lower-mantle, and were buried in epoxy mounts. XANES spectra were recorded in the fluorescence mode at the BL4A of the Photon Factory, KEK, Japan. Relative content of Cr²⁺ was calculated using the method described by Eeckhout et al. (2007).

The obtained results revealed that divalent chromium are present in the natural ferropericlasite and that the lower-mantle is reduced condition. The Cr²⁺ proportion to the total Cr content varied among the samples in the range of 3 to 10%. In contrast, no remarkable difference was observed in the valence state of Fe among the samples. Thus, the valence state of Cr in ferropericlasite can be a indicator of the redox environment in the lower-mantle.

キーワード: ダイヤモンド, 酸化還元状態, 下部マントル

Keywords: Diamond, Lower mantle