Japan Geoscience Union Meeting 2010

(May 23-28 2010 at Makuhari, Chiba, Japan)

©2009. Japan Geoscience Union. All Rights Reserved.



SGL045-P01

会場:コンベンションホール

時間: 5月25日17:15-18:45

四万十付加体の震源断層におけるフィッション・トラック分析

Zircon fission track thermochronology of an ancient seismogenic zone in Shimanto accretionary complex Southwest Japan

井上 文1, 郁芳 隨徹2*, 坂口 有人3, 田上 高広2, 渡邊 裕美子2

Aya Inoue¹, Zuitetsu Ikuho^{2*}, Arito Sakaguchi³, Takahiro Tagami², Yumiko Watanabe²

¹京都大学理学部理学科, ²京都大学大学院理学研究科, ³海洋研究開発機構

¹Kyoto University, ²Kyoto University, ³JAMSTEC

To better understand heat generation and transfer along faults in the subduction zone, we measured 7 length distributions of fission tracks in zircons separated from sandstones in and around ancient seismogenic zone found in the Shimanto accretionary complex, southwest Japan. Samples were collected along a traverse approximately orthogonal to the tectonic boundary fault between the Nonokawa Formation and Okitsu Melange of the complex, which was once located in the seismogenic zone judged from the occurrence of pseudotachylyte therin. The observed FTs showed significantly reduced mean length for samples from 10 to 20 meters away orthogonal to the fault on the hanging-wall side. This suggests that the heating up to the zircon partial annealing zone occurred by a thermal event along the deposition that locally perturbed the geothermal structure.

キーワード:断層,付加体,フィッション・トラック

Keywords: Fault, Acrretionary Complex, Fission Track