

放射光X線その場観察による高温高圧下における蛇紋石の状態方程式

Equations of state of antigorite under high pressure and high temperature determined by in situ X-ray diffraction (XRD)

Yang Cuiping^{2*}, 井上 徹¹, 金 振民², 山田 明寛¹, 曾我部 昭人¹, 松影 香子¹, 亀卦川卓美³

Cuiping Yang^{2*}, Toru Inoue¹, Zhenmin Jin², Akihiro Yamada¹, Akihito Sogabe¹,
Kyoko Matsukage¹, Takumi Kikegawa³

¹愛媛大学地球深部ダイナミクス研究センター, ²中国地質大学 (武漢), ³高エネルギー加速器研究機構

¹GRC, Ehime University, ²China University of Geosciences (Wuhan), ³PF, KEK

Many phase equilibrium experiments were conducted to clarify the dehydration phase boundary of antigorite; however, the results are inconsistent with each other. The unit-cell volume under high temperature and pressure (i.e. P-V-T equations of state (P-V-T EoS)) is one of the most important thermodynamic parameters of mineral to calculate the stability field using thermodynamic based calculation, in addition to discuss the density of serpentine-bearing subducted slab under high pressure and high temperature. But the existed EoS data of antigorite are few, and most of them were obtained by experiments at room temperature. Recently, Hilairet et al. (2006) conducted an in situ X-ray diffraction experiment under room temperature and high pressure to calculate the bulk modulus of antigorite. Their result is quite different from previous researches' (Bose and Navrotsky, 1998; Holland and Powell, 1998), so the stability field calculated by the thermodynamic is also different. The stability field of antigorite from Hilairet et al. (2006) is consistent with the result of the phase equilibrium experiment by Komabayashi et al. (2005). In this study, we have conducted in situ X-ray diffraction experiments under high pressure and high temperature up to about 8 GPa, 500 degree C to obtain the P-V-T EoS of antigorite, using 6-6 type multi-anvil (Nishiyama et al., 2008) in photon factory-advanced ring (PF-AR), Tsukuba, Japan. Our antigorite was stable during the whole pressure and temperature range in our experiments; the P-V-T data were fitted by 2nd order of Birch-Murnaghan EoS (fixed $K_0'=4$), and the obtained bulk modulus under different temperatures were: $K_{0,300K}=68.2(17)$, $K_{0,373K}=62.9(17)$, $K_{0,473K}=61.4(17)$, $K_{0,573K}=63.0(17)$, $K_{0,673K}=61.4(16)$, and $K_{0,773K}=61.6(25)$, respectively; the bulk modulus under room temperature was consistent with Hilairet et al. (2006). The c axis was more compressible compared to the other axes, a axis had the smallest compressibility, but the difference between a and b axes was small. The thermal expansion parameter of antigorite ($a_0=1.8 \text{ 0E-5/K}$) was comparable with chlorite (2.5E-5/K) (Pawley et al., 2002). Further details will be presented in the session.

Bose, K. and Navrotsky, A., 1998. Thermochemistry and phase equilibria of hydrous phases in the system MgO-SiO₂-H₂O: Implications for volatile transport to the mantle. *J. Geophys. Res.*, 103: 9713-9719.

Hilairet, N., Daniel, I. and Reynard, B., 2006. Equation of state of antigorite, stability field of serpentines, and seismicity in subduction zones. *Geophys. Res. Lett.*, 33: L02302.

Holland, T.J.B. and Powell, R., 1998. An internally consistent thermodynamic data set for phases of petrological interest. *Journal of Metamorphic Geology*, 16(3): 309-343.

Komabayashi, T., Hirose, K., Funakoshi, K.-i. and Takafuji, N., 2005. Stability of phase A in antigorite (serpentine) composition determined by in situ X-ray pressure observations. *Physics of The Earth and Planetary Interiors*, 151(3-4): 276-289.

Nishiyama, N., Wang, Y., Sanehira, T., Irifune, T. and Rivers, M.L., 2008. Development of the Multi-anvil Assembly 6-6 for DIA and D-DIA type high-pressure apparatuses. *High Pressure Research: An International Journal*, 28(3): 307 - 314.

Pawley, A.R., Clark, S.M. and Chinnery, N.J., 2002. Equation of state measurements of chlorite, pyrophyllite, and talc. *American Mineralogist*, 87(8-9): 1172-1182.

キーワード: 蛇紋石, アンチゴライト, 状態方程式, 放射光 X 線その場観察, 体積弾性率, 熱膨張率

Keywords: serpentine, antigorite, equation of state, in situ X-ray diffraction, bulk modulus, thermal expansion