## **Japan Geoscience Union Meeting 2010**

(May 23-28 2010 at Makuhari, Chiba, Japan)

©2009. Japan Geoscience Union. All Rights Reserved.



SRD051-P05

会場:コンベンションホール

時間: 5月23日17:15-18:45

フィリピン共和国、パラワン島、Rio Tuba、ラテライト型ニッケル鉱床におけるニッケルの地球化学的挙動

Geochemical behavior of nickel in weathered profile of the Rio Tuba laterite deposit, Palawan Island, Philippines

清水 公輔1\*, 林 謙一郎2

Kousuke Shimizu<sup>1\*</sup>, Ken-ichiro Hayashi<sup>2</sup>

<sup>1</sup>筑波大学大学院 生命環境科学研究科, <sup>2</sup>筑波大学大学院 生命環境科学研究科

<sup>1</sup>University of Tsukuba, <sup>2</sup>University of Tsukuba

Serpentinized ultramafic rocks accompanied by ophiolite are widely distributed at southern Palawan Island. Laterite type Ni deposit at the Rio Tuba mine is developed over the weathered serpentine body. It produces low grade Ni-ore, and after the introduction of high pressure acid leach (HPAL) technology, about 1 % of world nickel production comes from the deposit. Weathered profile consists of dark brown zone, reddish to brown zone and yellowish gray zone from surface to bottom. Because yellowish gray zone of >8 m from the surface poorly possesses original texture, this zone corresponds to saprolite. Upper zone of <8 m corresponds to laterite. Results of chemical analysis on bulk samples show that laterite zone is rich in Fe<sub>2</sub>O<sub>3</sub>(about 60 wt. %), and contains Al<sub>2</sub>O<sub>3</sub>(about 10 wt.%), SiO<sub>2</sub>(about 8 wt.%) and Cr<sub>2</sub>O<sub>3</sub>(about 5 wt.%). Iron content decreases in saprolite (Fe<sub>2</sub>O<sub>3</sub>= about 15 wt.%), while SiO<sub>2</sub>(about 35 wt.%) and MgO (about 25 wt. %) contents increase. Nickel is accompanied in saprolite with maximum content of 2.3 wt.%. XRD and IR-absorption studies suggest that laterite consists of hematite-goethite, and saprolite consists of upper smectite-talc and lower lizardite zones.

In order to determine host minerals of nickel, sequential extraction experiments have been conducted. Nickel is extracted from soil samples with reaction of various chemical regents. Nickel contents in fractions of 'solubl' 'exchangeable', 'carbonate phase', 'oxide phase', 'organic matter' and 'silicate and residual phases' will be separately obtained. Relationship between ore mineralogy and nickel content in each fraction will be discussed.

キーワード:ニッケル,ラテライト,連続抽出法,フィリピン

Keywords: nickel, laterite, sequential extraction, Philippines