

SSS019-03

会場: 303

時間: 5月23日16:00-16:15

海洋底堆積物中の擬弾性歪回復から明らかになった、西南日本沈み込み帯の広域的伸長応力場

Anelastic strain recovery in ocean floor sediments reveals extension across SW Japan subduction zone

Byrne Timothy B.^{1*}, 林 為人², 堤 昭人³, 山本 裕二⁷, Lewis Jonathan C.⁴, 金川 久一⁵, 北村 有迅⁶, 山口 飛鳥⁷, 木村 学⁸

Timothy B. Byrne^{1*}, Weiren Lin², Akito Tsutsumi³, Yuhji Yamamoto⁷, Jonathan C. Lewis⁴, Kyuichi Kanagawa⁵, Yujin Kitamura⁶, Asuka Yamaguchi⁷, Gaku Kimura⁸

¹コネチカット大学, ²海洋研究開発機構, ³京都大学, ⁴ペンシルバニアインディアナ大学, ⁵千葉大学, ⁶キール大学, ⁷高知大学, ⁸東京大学

¹University of Connecticut, ²JAMSTEC, ³Kyoto University, ⁴Indiana University of Pennsylvania, ⁵Chiba University, ⁶University of Kiel, ⁷Kochi University, ⁸University of Tokyo

Sediment dominated convergent margins typically record substantial horizontal shortening often associated with great earthquakes. The convergent margin south of Japan is arguably one of the most extensively investigated margins, and previous studies have documented extensive evidence for accretion and horizontal shortening. Here, we show results from anelastic strains recovered from three partially lithified sediment samples (ca 40% porosities) across the southwest Japan accretionary prism and propose that the shallow structural levels of the margin (to at least 900 m) are dominated by horizontal extension rather than compression. The anelastic strain results are consistent with stress directions interpreted from two independent techniques - bore hole breakout orientations and core-scale fault data. The anelastic strain axes are also oblique to bedding and, at one site, oblique to fabrics defined by AMS. The AMS fabrics dip steeply, consistent with horizontal tectonic stresses and shortening. Based on the pervasive occurrence of late-stage normal faults and the ASR data, we interpret the AMS data to reflect remnant fabrics from an earlier phase of shortening. We propose that the current state of stress reflects the relatively recent (<1 million years) underthrusting of a thick sediment pile that increased the wedge taper and weakened the decollement.

キーワード: 南海トラフ沈み込み帯, 擬弾性歪回復, 孔内破壊, 小断層解析, 帯磁率異方性, 伸長応力場

Keywords: Nankai Trough subduction zone, anelastic strain recovery, borehole breakout, fault-slip analysis, anisotropy of magnetic susceptibility, extension