

IODP第322次航海の初期成果：南海トラフ沈み込みインプット

Preliminary Results from IODP Expedition 322: NanTroSEIZE Subduction Inputs

斎藤 実篤^{1*}, マイク・アンダーウッド², 久保雄介³, 真田佳典³, 荒木英一郎⁴,
ティム・バーン⁵, リサ・マクニール⁶, デミアン・セイファー⁷, 江口暢久³, 高橋共馬³,
ショーン・トシコ³, モー・キョー・トゥ³, IODP第319/322次研究航海乗船研究者³

Saneatsu Saito^{1*}, Michael B. Underwood², Yusuke Kubo³, Yoshinori Sanada³, Eiichiro Araki⁴,
Tim Byrne⁵, Lisa McNeill⁶, Demian Saffer⁷, Nobuhisa Eguchi³, Kyoma Takahashi³,
Sean Toczko³, Moe Kyaw Thu³, IODP Expedition 319/322 Scientists³

¹海洋研究開発機構地球内部ダイナミクス領域, ²ミズーリ大学, ³海洋研究開発機構地球深部探査センター,
⁴海洋研究開発機構地震津波・防災研究P, ⁵コネチカット大学, ⁶サザンプトン大学, ⁷ペンシルバニア大学

¹IFREE, JAMSTEC, ²University of Missouri, ³CDEX, JAMSTEC, ⁴DONET, JAMSTEC, ⁵University of Connecticut,
⁶University of Southampton, ⁷Pennsylvania State University

Integrated Ocean Drilling Program (IODP) Expedition 322 was designed to document characteristics of incoming sedimentary strata and igneous basement prior to their arrival at the subduction front. To accomplish these objectives, coring was conducted at two sites in the Shikoku Basin on the subducting Philippine Sea plate. Site C0011 is located on the northwest flank of the Kashinosaki Knoll, whereas Site C0012 is located near the crest of the knoll. The resulting data, which include logging-while-drilling during IODP Expedition 319, provide a wealth of new information on presubduction equivalents of the seismogenic zone. Near the end of Expedition 319, LWD data were collected at proposed Site NT1-07 (Hole C0011A). LWD data include natural gamma radiation, resistivity, and resistivity images. We were able to make confident correlations and comparisons between the logs and subsequent core description and multisensor core logger (MSCL) data. The whole section was divided into five logging units on the basis of visual inspection of the log responses. These divisions also correlate reasonably well with the seismic stratigraphy. Analysis of borehole breakouts indicates that the maximum horizontal stress field (SHmax) is orientated north-northeast-south-southwest, roughly perpendicular to the convergence direction of the Philippine Sea plate. Although coring at Site C0011 failed to reach the total depth target, coring at Site C0012 penetrated 33 m into igneous basement and recovered the sediment/basalt interface intact at ~540 m CSF. The age of basal sediment is >18.9 Ma. The merger of lithofacies and age-depth models from the two sites captures all of the important ingredients of basin evolution; including an unrecognized interval of late Miocene tuffaceous/volcaniclastic sandstone designated the middle Shikoku Basin facies. An older (early to middle Miocene) turbidite sandstone/siltstone facies with mixed detrital provenance occurs in the lower Shikoku Basin; this unit is broadly correlative with similar Miocene turbidites on the western side of the basin. Another result came from geochemical analyses of pore water and hydrocarbons at Site C0012. Pore fluids on top of the basement high show clear evidence of a seawater-like source, with chlorinity values increasing toward basement because of hydration reactions and diffusion; the fluids are largely unchanged by the effects of focused flow and/or in situ dehydration reactions associated with rapid burial beneath the trench wedge and frontal accretionary prism. Thus, Site C0012 finally provides a reliable geochemical reference site for the subduction zone.

Keywords: Integrated Ocean Drilling Program, NanTroSEIZE, subduction inputs, D/V Chikyu, Expedition 322, logging-while-drilling