

SSS020-06

会場: 302

時間: 5月26日10:15-10:30

コディアク付加体Pasagshak Point thrustにおける地震時の水-岩石相互作用

Co-seismic water-rock interaction recorded in Pasagshak Point thrust, Kodiak accretionary complex, Alaska

山口 飛鳥^{1*}, 石川 剛志², 加藤 泰浩³, Moore J. C.⁴, Rowe Christie D.⁵, Meneghini F.⁶, 堤 昭人⁷, 氏家 恒太郎⁸, 木村 学⁹

Asuka Yamaguchi^{1*}, Tsuyoshi Ishikawa², Yasuhiro Kato³, J. C. Moore⁴, Christie D. Rowe⁵, F. Meneghini⁶, Akito Tsutsumi⁷, Kohtaro Ujiie⁸, Gaku Kimura⁹

¹高知大学海洋コア総合研究センター, ²海洋研究開発機構高知コア研究所, ³東京大学大学院工学系研究科, ⁴カリフォルニア大学サンタクルズ校, ⁵ケープタウン大学, ⁶ピサ大学, ⁷京都大学大学院理学研究科, ⁸海洋研究開発機構地球内部ダイナミクス領域, ⁹東京大学大学院理学系研究科

¹Kochi Univ., ²JAMSTEC, ³Univ. Tokyo, ⁴UCSC, ⁵Univ. Cape Town, ⁶Univ. Pisa, ⁷Kyoto Univ., ⁸JAMSTEC, ⁹Univ. Tokyo

A pseudotachylite-like ultra-fine grained black fault rock (BFR) has discovered from Pasagshak Point thrust, Ghost Rocks formation, Kodiak Island, Alaska (Rowe et al., 2005). It is very different from pseudotachylites in Shimanto accretionary complex (Ikesawa et al., 2003; Kitamura et al., 2005; Okamoto et al., 2006; Mukoyoshi et al., 2006; Ujiie et al., 2007) in the aspect of its occurrences: i.e. large thickness (up to 20cm), lack of mineral veins, variety of mixing pattern with surrounding random and foliated cataclasite. Rowe et al. (submitted) and Meneghini et al. (2010) have explored meso-scale deformational features and microfabrics, respectively, and concluded that the BFR had been formed by ancient seismic events occurred in plate boundary decollement at seismogenic depths (temperatures approximately 240 to 260 C).

In this study, we have performed analyses of whole-rock trace element concentrations and ⁸⁷Sr/⁸⁶Sr ratios of BFRs, cataclasites, and host rocks on the basis of detailed structural analyses.

Enrichment in Sr, depletions in Rb and Cs in comparison with host rocks suggest that the BFR have suffered high-temperature water-rock interactions larger than 350 C (Ishikawa et al., 2008). Positive Li anomaly and low ⁸⁷Sr/⁸⁶Sr ratio of BFR imply that surrounding fluid was enriched in Li and ⁸⁶Sr. Rb-Sr isochron plot might suggest that Rb-Sr age was reset due to mechanochemical generation of plagioclase microcrystals. Intense water-rock interactions in spite of the lack of mineral veins would affect sealing and earthquake preparation processes in seismic cycles. Kodiak BFRs present new pattern of water-rock interaction at seismogenic depth of subduction zones.

Keywords: Kodiak accretionary complex, decollement, water-rock interaction, seismogenic zone, earthquake, fault rock