

SSS025-13

会場: 301A

時間:5月25日15:30-15:45

ALOS/PALSARデータから推定された文川地震における断層傾斜角の変化

Spatial changes of the fault dip angle in the Wenchuan earthquake, China, as inferred from ALOS/PALSAR data

古屋 正人1*, 小林 知勝², 高田 陽一郎³, 村上 亮1

Masato Furuya^{1*}, Tomokazu Kobayashi², Youichiro Takada³, Makoto Murakami¹

¹北海道大学理学研究院,²国土地理院,³海洋研究開発機構

¹Hokkaido University, ²Geographical Survey Institute, ³JAMSTEC

We use crustal deformation data sets derived from satellite-based synthetic aperture radar (SAR) to develop a fault source model of the 2008 M7.9 Wenchuan earthquake, China, which occurred at the Longmen Shan Fault Zone. The data sets include interferometric SAR (InSAR), range offset, and azimuth offset data acquired at seven ascending paths. The range and azimuth offset data are particularly important, exhibiting a single major rupture to the northeast (NE) and multiple ruptures to the southwest (SW). Our preferred model consists of six segments; four follow the previously mapped traces of the Beichuan fault (BF) and its NE extension, one corresponds to the Pengguan fault (PF) to the SW, and the other is included to represent a conjugate fault to the SW. Maximum slip of $^{-10}$ m consisting of both thrust and right-lateral slip components is identified at the shallowest patches to the NE along the BF. The multiple fault segments to the SW show that the thrust slip component initially dominates, and the strike slip becomes significant toward the NE. Examining the sensitivity of the predicted azimuth offset data to the assumed dip angle, we found that the dip angle changed significantly from 35–45 degrees at the SW fault segment for the PF to 80--90 degrees at the NE extension of the BF. Without using ground-based data, we have independently confirmed that the fault dip angle evolved from the SW to the NE, which is reported by Shen et al. (2009).

キーワード:四川地震,断層傾斜角,合成開口レーダー,地殻変動

Keywords: Wenchuan earthquake, Fault dip angle, SAR, crustal deformation, ALOS, PALSAR