

AAS021-01

会場:102

時間:5月23日 08:30-08:45

## 地上観測および CONTRAIL による大気 CO<sub>2</sub> データを用いた炭素収支の逆解析 Carbon budget estimation by inverse modeling with atmospheric CO<sub>2</sub> concentrations from surface and CONTRAIL measurements

丹羽 洋介<sup>1\*</sup>, 町田 敏暢<sup>2</sup>, 澤 庸介<sup>1</sup>, 松枝 秀和<sup>1</sup>, 今須 良一<sup>3</sup>, 佐藤 正樹<sup>3</sup>

Yosuke Niwa<sup>1\*</sup>, Toshinobu Machida<sup>2</sup>, Yousuke Sawa<sup>1</sup>, Hidekazu Matsueda<sup>1</sup>, Ryoichi Imasu<sup>3</sup>, Masaki Satoh<sup>3</sup>

<sup>1</sup> 気象研究所, <sup>2</sup> 国立環境研究所, <sup>3</sup> 東京大学大気海洋研究所

<sup>1</sup>MRI, <sup>2</sup>NIES, <sup>3</sup>AORI

A great deal of understanding of the global and regional carbon budget helps us to perform a reliable prediction of future climate with an earth system model. However, the accuracy of CO<sub>2</sub> source/sink estimation by inverse modeling, which is one of the leading methods to estimate regional carbon budget, is not very high because of sparse observational data coverage. The recent evolving aircraft measurements of CO<sub>2</sub> in a three-dimensional view are expected to provide new constraints on the estimation of surface CO<sub>2</sub> fluxes.

In this study, regionally divided carbon budgets are estimated by inverse modeling using surface measurement networks and aircraft measurements from Comprehensive Observation Network for Trace gases by Airliner (CONTRAIL). The CONTRAIL project has started since late 2005 and a huge amount of atmospheric CO<sub>2</sub> data has been obtained covering altitudes between the Earth surface to the upper-troposphere and lower-stratosphere, latitudes between the boreal high-latitudes to the austral mid-latitudes. Monthly mean observational data from GLOBALVIEW-CO<sub>2</sub>, which mostly consists of surface measurements, and CONTRAIL are used in this inverse analysis. The CONTRAIL data measured both vertically over each airport and horizontally at the cruising altitude are used. The inversion method is based on the Bayesian statistics and the approach of the TransCom 3 is used. To relate atmospheric concentrations to surface fluxes, a three-dimensional transport model is employed. In this study, Nonhydrostatic ICosahedral Atmosphere Model (NICAM)-based transport model (NICAM-TM) is used with prescribed CO<sub>2</sub> flux data of fossil fuel emission, respiration/photosynthesis in terrestrial biosphere and atmosphere-ocean exchange. The transport simulations of atmospheric CO<sub>2</sub> are performed by low-resolution version of NICAM; the horizontal grid interval is about 240 km. The analyzed period is five years during 2005-2009 and meteorological fields in each year are used to drive the transport model with the nudging method. The inversion setup is similar to TransCom3, but number of flux regions to be estimated is 42. Land regions are divided into 31 according to vegetation types and the same 11 ocean regions as TransCom 3 are used.

A preliminary result of forward simulation with the prescribed fluxes shows that the transport model has good performance for reproducing general features of three-dimensional structure of CO<sub>2</sub> observed by CONTRAIL. However, some discrepancies between the simulation and CONTRAIL are found in horizontal gradient even in the upper-troposphere during summer, indicating that aircraft measurements have significant impacts on flux estimates when vertical transport is efficient. The inversion results will be discussed in the presentation.

キーワード: 炭素循環, 逆解析

Keywords: carbon cycle, inversion