

AAS021-02

会場:102

時間:5月23日 08:45-09:00

## CONTRAIL 輸送モデル比較実験で再現された上部対流圏/下部成層圏のCO<sub>2</sub>濃度分布の季節変化 CO<sub>2</sub> seasonal distributions in the UT/LS region as observed by CONTRAIL and four transport models

澤 庸介<sup>1\*</sup>, 丹羽洋介<sup>1</sup>, Prabir K. Patra<sup>2</sup>, 町田敏暢<sup>3</sup>, 松枝秀和<sup>1</sup>, Dmitry Belikov<sup>3</sup>, 池上雅明<sup>4</sup>, 眞木貴史<sup>1</sup>, Shamil Maksyutov<sup>3</sup>, 小田知宏<sup>3</sup>, 今須良一<sup>5</sup>, 佐藤正樹<sup>5</sup>

Yousuke Sawa<sup>1\*</sup>, Yosuke Niwa<sup>1</sup>, Prabir K. Patra<sup>2</sup>, Toshinobu Machida<sup>3</sup>, Hidekazu Matsueda<sup>1</sup>, Dmitry Belikov<sup>3</sup>, Masaaki Ikegami<sup>4</sup>, Takashi Maki<sup>1</sup>, Shamil Maksyutov<sup>3</sup>, Tomohiro Oda<sup>3</sup>, Ryoichi Imasu<sup>5</sup>, Masaki Satoh<sup>5</sup>

<sup>1</sup> 気象研究所, <sup>2</sup> 海洋研究開発機構, <sup>3</sup> 国立環境研究所, <sup>4</sup> 気象庁環境気象管理官, <sup>5</sup> 東京大学大気海洋研究所

<sup>1</sup>Meteorological Research Institute, <sup>2</sup>JAMSTEC, <sup>3</sup>National Inst. for Environmental Studies, <sup>4</sup>Japan Meteorological Agency,

<sup>5</sup>AORI, The University of Tokyo

In the Comprehensive Observation Network for Trace gases by AirLiner (CONTRAIL) project, high-frequency and wide-ranging CO<sub>2</sub> data in the upper troposphere(UT)/lower stratosphere(LS) region have been obtained by Continuous CO<sub>2</sub> Measuring Equipment (CME) onboard commercial aircraft operated by Japan Airlines (JAL).

The observed distributions in UT/LS region showed that CO<sub>2</sub> isopleths followed the tropopause during the winter and spring. On the other hand, distributions tracked potential temperature surfaces crossing the tropopause in summer, suggesting fast meridional transport of high CO<sub>2</sub> from the tropical troposphere. However, it is difficult to show how the tropospheric air masses intrude into the lower stratosphere across the tropopause due to the limited observational data.

Recently, we conducted CONTRAIL transport model intercomparison to improve our knowledge of three-dimensional structures of atmospheric CO<sub>2</sub>. The distributions in UT/LS region simulated by four global chemical transport models (ACTM, MJ98-CDTM, NICAM-TM, NIES) with common CO<sub>2</sub> flux dataset were used to study the transport processes near the tropopause. It was shown that the models reproduced observed CO<sub>2</sub> distributions following the tropopause in winter-spring season, although the CO<sub>2</sub> gradients across the tropopause are underestimated by approximately 2 ppm between 300 and 350 K in potential temperature surfaces. In summer, the isentropic transport of high CO<sub>2</sub> from the upper-troposphere in lower latitudes to the lower-stratosphere in higher latitudes were well simulated by the models. The detailed processes of transport will be analyzed and shown in the meeting.

キーワード: 二酸化炭素, 輸送モデル, 対流圏成層圏交換

Keywords: CO<sub>2</sub>, transport model, UT/LS exchange