

AAS021-13

会場:102

時間:5月23日 11:45-12:00

2010年春季における福江島および中国如東でのPM2.5濃度変動 PM2.5 variation in Rudong, China and Fukue Island, Japan in spring 2010

竹谷 文一^{1*}, 金谷 有剛¹, Xiaole Pan¹, 高島 久洋¹, 入江 仁士¹, 駒崎雄一¹, 斉藤伸治², 猪俣 敏², 谷本 浩志², Huabin Dong³, Baozhu Ge³, Jie Li³, Xiquan Wang³, Hongyan Chen³, Meigen Zhang³, Zifa Wang³
Fumikazu Taketani^{1*}, Yugo Kanaya¹, Xiaole Pan¹, Hisahiro Takashima¹, Hitoshi Irie¹, Yuichi Komazaki¹, Shinji Saito², Satoshi Inomata², Hiroshi Tanimoto², Huabin Dong³, Baozhu Ge³, Jie Li³, Xiquan Wang³, Hongyan Chen³, Meigen Zhang³, Zifa Wang³

¹ 海洋研究開発機構, ² 国立環境研究所, ³ 中国科学院大气物理研究所

¹JAMSTEC, ²NIES, ³IAP

We carried out an intensive field campaign observing ozone and its precursors and chemical components/physical and optical parameters of aerosol particles at Rudong (32.26N, 121.37E), Jiangsu, China in May/June 2010. In this presentation, to investigate the PM2.5 mass concentration, their composition and correlation in both Rudong and Fukue Island(32.75N, 128.68E), we observed PM2.5 mass concentration using SHARP monitor and sampled PM2.5 on the quartz filter by high volume air sampler. PM2.5 total mass concentration is monitored every 1 min for both sites. At the Rudong site(May16-June23), 9 or 14-hours PM2.5 samples were collected on the quartz filters using High-volume(500L/min) sampler, while 22-hours PM2.5 samples were collected at the Fukue Island site(May18-25). In the end of May, high PM2.5 mass concentrations were observed in both Rudong and Fukue Island. We are going to discuss mass closure and correlation of the PM2.5 for Rudong and Fukue Island results.

Keywords: PM2.5, aerosol, composition, transportation, mass closure