

ACG004-12

会場:104

時間:5月27日 09:15-09:30

熱帯域雲活動への太陽活動とQBOの影響

Implications for the low latitude cloud formations from solar activity and the quasi-biennial oscillation

宮原 ひろ子^{1*}, 洪 鵬², 横山 祐典², 高橋 幸弘³, 佐藤 光輝³
Hiroko Miyahara^{1*}, Peng Hong², Yusuke Yokoyama², Yukihiro Takahashi³, Mitsuteru Sato³

¹ 東京大学宇宙線研究所, ² 東京大学, ³ 北海道大学

¹ICRR, The University of Tokyo, ²The University of Tokyo, ³Hokkaido University

We examined the effect of the 11-year solar cycle and quasi-biennial oscillation (QBO) on the 27-day solar rotational period detected in tropical convective cloud activity. We analyzed the data of outgoing longwave radiation (OLR) for AD1979-2004, dividing into four different cases by the combination of high and low solar activities in terms of the 11-year variation, and easterly and westerly stratospheric winds associated with QBO. As a result, 27-day variation has been most significantly detected in high solar activity period around the Indo-Pacific Warm Pool. Based on correlation analysis, we find that solar rotation signal can explain 10-20% of OLR variability around the tropical warm pool region during the high solar activity period. The spatial distribution has been, however, apparently different according to the phases of QBO. It is suggested that the 11-year solar cycle and stratospheric QBO have a possibility to cause large-scale oceanic dipole phenomena.

キーワード: 太陽活動, 成層圏準2年周期振動 (QBO), 西太平洋域, 雲活動

Keywords: Outgoing Longwave Radiation (OLR), solar activity, solar rotation, Quasi Biennial Oscillation (QBO), Western Pacific Region