## **Japan Geoscience Union Meeting 2011**

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



ACG032-22 会場:105

時間:5月27日15:30-15:45

## 20世紀前半の西部北太平洋域の台風経路データセット Historical typhoon track dataset during the early 20th century over the western north Pacific

久保田 尚之 1\*, 松本 淳 2, Edwin Ginn<sup>3</sup> Hisayuki Kubota 1\*, Jun Matsumoto<sup>2</sup>, Edwin Ginn<sup>3</sup>

1 海洋研究開発機構, 2 首都大学東京, 3 香港気象局

Typhoon provides fresh water to the land but it can cause disaster when it makes landfall due to strong winds and heavy rain. Recently the variability of typhoon activity becomes a great concern because it may be affected by global warning. Over the Western North Pacific (WNP) basin, typhoon best track data are available from 1945. Before that due to the difference of typhoon definition, there is no available database. In this study, we collected and digitized the historical typhoon track data during the early 20th century. We created new typhoon definition, performed quality check by comparing station pressure data, connected to the current best track data and made 100 years typhoon track dataset over the western north Pacific. The purpose of this study is to understand the typhoon variability during the 20th century. Now we have four different sources of historical typhoon track data. By comparing different sources we are improving the typhoon track data.

It is important to keep the quality of the historical typhoon track data as close as current best track data to make the dataset reliable. We would like to discuss how to keep this quality of the dataset and the availability of the dataset to other communities.

キーワード: 台風, 西部北太平洋, データレスキュー, 気候変動

Keywords: typhoon, western north Pacific, data rescue, climate change

<sup>&</sup>lt;sup>1</sup>JAMSTEC, <sup>2</sup>Tokyo Metropolitan University, <sup>3</sup>Hong Kong Observatory