

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



ACG034-P08

会場:コンベンションホール

時間:5月27日 14:00-16:30

全球風化速度の定量化に向けた化学風化モデルの開発

Development of a chemical weathering model toward quantification of global weathering rate in the past

野津 太一^{1*}, 田近 英一¹

Taichi Nozu^{1*}, Eiichi Tajika¹

¹ 東京大学理学系研究科地球惑星科学専攻

¹Dept. EPS, Sch. Sci., Univ. Tokyo

Global climate is thought to have been maintained by the long-term balance of CO₂ due to continental silicate weathering and volcanic degassing. The rate of chemical weathering of silicate minerals depends on lithology, temperature, runoff, plant evolution, soil microbial activity, and so on. Although the relationship among the controlling factors of the chemical weathering rate is still an open question, many studies have mentioned the importance of the rate of physical weathering, or erosion, on the chemical weathering rate. Efforts to quantify the chemical weathering rate using numerical models have also been made, but the number of publications in which such kind of process-based models were applied to paleo-environmental study is limited. We try to develop a process-based weathering model which can be applied for investigating roles of chemical weathering in paleo-environmental change through biogeochemical cycle quantitatively on the basis of governing physical, chemical, and biological processes.

キーワード: 化学風化, 数値モデル

Keywords: chemical weathering, process-based modeling