## **Japan Geoscience Union Meeting 2011**

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



AHW026-01 Room:201A Time:May 27 14:15-14:30

## Distribution properties of phosphorus concentration in coastal groundwater: hydrogeological background

Shin-ichi Onodera<sup>1\*</sup>, Mitsuyo Saito<sup>2</sup>, Mayumi Jige<sup>3</sup>, Yuta Shimizu<sup>1</sup>, Guangzhe Jin<sup>1</sup>, Seiko Yoshikawa<sup>4</sup>

<sup>1</sup>Hiroshima University, <sup>2</sup>Ehime University, <sup>3</sup>Chiba Science University, <sup>4</sup>Research Institute of Agriculture

Nutrient condition in water environment controls the ecosystem. Groundwater discharge to the oceans is significant as nutrient supply (Slomp et al, 2004 etc). Especially, phosphorus and silica concentration generally are relatively high in coastal area. However, it has not been enough to confirm the source of phosphorus in coastal groundwater in previous studies. This study aims to confirm hydrogeological properties in coastal groundwaters, and estimate the possible phosphorus sources of groundwater.

The study areas are Osaka, Marugame, Okayama, and Fukuyama alluvial plains and small island groundwaters in Hiroshima prefecture. We arranged hydrogeological and groundwater quality data sets in previous studies of Hiroshima University. The phosphorus concentrations were high in anoxic condition. In addition, shallow aquifers around alluvial clay had high concentrations. The phosphorus contents in alluvial sediments of Okayama plain were relatively high around alluvial clay. These results suggest the contribution of phosphorus from alluvial sediment to groundwater.

Keywords: phosphorus, coastal groundwater, hydrogeology, sediment